

## STUDIES IN VIBRATIONAL SPECTROSCOPY

- I. NORMAL COORDINATE ANALYSIS OF  $\text{CF}_3\text{OF}$  AND  $\text{CF}_3\text{OCF}_3$
- II. RAMAN SPECTRA OF  $\text{CF}_2(\text{OF})_2$  AS A FUNCTION OF TEMPERATURE

by

JENG-CHUNG KUO

B.S., National Taiwan University, 1972

---

A MASTER'S THESIS

Submitted in partial fulfillment of the  
requirements for the degree

MASTER OF SCIENCE

Department of Chemistry

KANSAS STATE UNIVERSITY  
Manhattan, Kansas

1978

Approved by:

R. M. Hammaker  
Major Professor

Document  
LD  
2668  
.T4  
1978  
K86  
C.2

TABLE OF CONTENTS

	Page
LIST OF TABLES	iv
LIST OF FIGURES	vi
PART I	
NORMAL COORDINATE ANALYSIS OF $\text{CF}_3\text{OF}$ AND $\text{CF}_3\text{OCl}$	
CHAPTER I. INTRODUCTION AND STATEMENT OF PROBLEM	1
CHAPTER II. EXPERIMENTAL SECTION	6
A. Recording of Spectra	6
1. Materials	6
2. Infrared Instrumentation	6
3. Raman Instrumentation	6
B. Computer Programs	7
CHAPTER III. EXPERIMENTAL RESULTS	9
A. Infrared Spectra of $\text{CF}_3\text{OF}$	9
B. Raman Spectra of $\text{CF}_3\text{OF}$	9
C. Infrared Spectra of $\text{CF}_3\text{OCl}$	10
D. Raman Spectra of $\text{CF}_3\text{OCl}$	10
E. Matrix Spectra of Smardzewski and Fox	10
CHAPTER IV. NORMAL COORDINATE ANALYSIS	23
A. Structural Models	23
B. Construction of the G Matrix	24
C. Utilization of Symmetry	27
D. Choice of Force Fields	27

	Page
E. Chemical Approach <u>vs.</u> Mathematical Approach	28
CHAPTER V. RESULTS, CONCLUSIONS AND OTHER REMARKS	
A. Comments on the Choice of Force Fields	37
B. Comments on Assignments Implied by Force Constant Calculations	41
C. Comments on the $L^{-1}$ Matrices and the Form of the Normal Coordinates	54
D. Comments on the Potential Energy Distribution	57
E. Comments on the Barrier to Internal Rotation	60

## PART II

RAMAN SPECTRA OF $CF_2(OF)_2$ AS A FUNCTION OF TEMPERATURE	
CHAPTER I. INTRODUCTION	64
CHAPTER II. EXPERIMENTAL	67
CHAPTER III. RESULTS AND DISCUSSION	72
A. General	72
B. Intensity Considerations	73
C. Energy Difference between Rotational Isomers	83
D. Suggestions for Future Work	85
LITERATURE CITED	87
ACKNOWLEDGMENT	91
VITA	92

## LIST OF TABLES

## PART I

NORMAL COORDINATE ANALYSIS OF  $\text{CF}_3\text{OF}$  AND  $\text{CF}_3\text{OCl}$ 

Table	Page
1. Raman and Infrared Data for $\text{CF}_3\text{OF}$ below $1400\text{ cm}^{-1}$	12
2. Raman and Infrared Data for $\text{CF}_3\text{OCl}$ below $1400\text{ cm}^{-1}$	17
3. Structural Data for $\text{CF}_3\text{OX}$ Compounds	25
4. Symmetry Coordinates for $\text{CF}_3\text{OX}$ Molecules of $\text{C}_s$ Symmetry	31
5. Internal Coordinate Force Constants for $\text{CF}_3\text{OF}$ and $\text{CF}_3\text{OCl}$	34
6. Assignments of the Fundamental Vibrations of Trifluoromethyl hypofluorite ( $\text{CF}_3\text{OF}$ )	35
7. Assignments of the Fundamental Vibrations of Trifluoromethyl hypochlorite ( $\text{CF}_3\text{OCl}$ )	36
8. Bond-stretching Force Constants of O-F and O-Cl Bonds from related Molecules	40
9. Frequency Comparison for Bands in the $200\text{-}300\text{ cm}^{-1}$ Region for $\text{CF}_3\text{OX}$ Compounds	44
10. Symmetrized $\text{L}^{-1}$ Matrix for $\text{CF}_3\text{OF}$	55
11. Symmetrized $\text{L}^{-1}$ Matrix for $\text{CF}_3\text{OCl}$	56
12. Potential Energy Distribution for $\text{CF}_3\text{OF}$	58
13. Potential Energy Distribution for $\text{CF}_3\text{OCl}$	59
14. Summary of Parameters in Barrier Height Calculation	63

Table

Page

## PART II

RAMAN SPECTRA OF  $\text{CF}_2(\text{OF})_2$  AS A FUNCTION OF TEMPERATURE

- |   |    |
|---|----|
| 15. Relative Intensity of the Bands from Liquid Raman Spectra of $\text{CF}_2(\text{OF})_2$ | 78 |
| 16. Summary of Tentative Assignment for a 2 Conformer Mixture                               | 82 |

## LIST OF FIGURES

## PART I

NORMAL COORDINATE ANALYSIS OF  $\text{CF}_3\text{OF}$  AND  $\text{CF}_3\text{OCl}$ 

Figure	Page
1. Gas-phase Infrared Spectra of $\text{CF}_3\text{OF}$ in the Region 1400-500 $\text{cm}^{-1}$	13
2. Gas-phase Infrared Spectra of $\text{CF}_3\text{OF}$ in the Region 650-200 $\text{cm}^{-1}$	14
3A. Raman Spectrum of Liquid $\text{CF}_3\text{OF}$	15
3B. Raman Spectrum of Gaseous $\text{CF}_3\text{OF}$	15
4. High Sensitivity Raman Spectra of Liquid $\text{CF}_3\text{OF}$	16
5. Gas-phase Infrared Spectra of $\text{CF}_3\text{OCl}$ in the Region 1400-500 $\text{cm}^{-1}$	18
6. Gas-phase Infrared Spectra of $\text{CF}_3\text{OCl}$ in the Region 650-180 $\text{cm}^{-1}$	19
7. Raman Spectrum of Liquid $\text{CF}_3\text{OCl}$	20
8. High Sensitivity Raman Spectra of Liquid $\text{CF}_3\text{OCl}$	21
9. $\nu(\text{OX})$ and $\nu(\text{CO})$ Stretches for $\text{CF}_3\text{OX}$ Molecules	22
10. Structure of $\text{CF}_3\text{OX}$ Molecules	26
11. Internal Coordinates for $\text{CF}_3\text{OX}$ Molecules	32
12. Unsymmetrized F Matrix for $\text{CF}_3\text{OX}$ Molecules	33

## PART II

RAMAN SPECTRA OF  $\text{CF}_2(\text{OF})_2$  AS A FUNCTION OF TEMPERATURE

13. Gas-phase Infrared Spectrum of $\text{CF}_2(\text{OF})_2$ in the Region 3000-300 $\text{cm}^{-1}$	69
14. Gas-phase Infrared Spectrum of $\text{CF}_2(\text{OF})_2$ in the Region 1400-1100 $\text{cm}^{-1}$	70

15. Raman Spectrum of Gas-phase $\text{CF}_2(\text{OF})_2$ in the Region 1000-800 $\text{cm}^{-1}$	71
16. Raman Spectrum of Liquid $\text{CF}_2(\text{OF})_2$ ( $-196^\circ\text{C}$ )	74
17. Raman Spectrum of Liquid $\text{CF}_2(\text{OF})_2$ ( $-135^\circ\text{C}$ )	75
18. Raman Spectrum of Liquid $\text{CF}_2(\text{OF})_2$ ( $-78^\circ\text{C}$ )	76
19. Raman Spectrum of Liquid $\text{CF}_2(\text{OF})_2$ ( $-40^\circ\text{C}$ )	77
20. Effect of Temperature on the Raman Spectrum of Liquid $\text{CF}_2(\text{OF})_2$ in the Range 1000-900 $\text{cm}^{-1}$	80
21. Effect of Temperature on the Raman Spectrum of Liquid $\text{CF}_2(\text{OF})_2$ in the Range 900-800 $\text{cm}^{-1}$	81

# I. NORMAL COORDINATE ANALYSIS OF $\text{CF}_3\text{OF}$ and $\text{CF}_3\text{OCl}$

## CHAPTER I INTRODUCTION AND STATEMENT OF PROBLEM

Normal coordinate analyses relate the observed, or preferably the harmonic, infrared and Raman vibrational frequencies to the force constants, equilibrium geometry, and atomic masses of the oscillating system (1). The development of increasingly reliable intramolecular force fields for a wide variety of molecules suggests that judicious applications of normal coordinate analyses may serve as an additional means for confidently probing the structural and bonding characteristics of diverse chemical systems.

For interpreting the vibrational spectra of larger systems, the concepts of modified potential functions and of the transferability of selected force constants from chemically similar but smaller molecules emerge.

The present study consists of two separate parts. The first part is a detailed study of the normal coordinate analyses of  $\text{CF}_3\text{OX}$  ( $\text{X}=\text{F}, \text{Cl}$ ) compounds. These two compounds are of interest because they are the initial members of the respective  $\text{R}_\text{f}\text{OF}$  and  $\text{R}_\text{f}\text{OCl}$  series, where  $\text{R}_\text{f}$  denotes the perfluoroalkyl group (2,3). The second part deals with the Raman spectrum of  $\text{CF}_2(\text{OF})_2$  and is discussed later in the thesis.

Our purpose in this part is twofold. The first objective



is to evaluate some results of previous vibrational analyses (4,5). With the complete analyses of the  $\text{CF}_3\text{OX}$  series, we will be able to see whether or not the vibrational frequencies chosen for fundamentals can be fitted reasonably well with a realistic potential function. Success in fitting the assignment with a realistic potential function will serve as evidence for the reliability of the new assignments which in turn may clarify some questionable results reported by Wilt and Jones (4) and support the work of Smardzewski and Fox (5) and of Buckley and Weber (6).

The second objective is to use the result of the  $\text{CF}_3\text{OX}$  series as the basis for later normal coordinate analyses of the  $\text{CF}_3\text{OOX}$  ( $\text{X}=\text{H}, \text{D}, \text{F}$  and  $\text{Cl}$ ) series, where electron diffraction data are now available (7), and to assist in the assignment of  $\text{CF}_3\text{OOX}$  series members which are isoelectronic with  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  (8).

In the present investigation, the calculations were carried out using the Wilson FG matrix method (9) with the computer programs written by Schachtschneider (10). Frequencies were given the weight  $1/(\lambda_i)^2$ , where  $\lambda = 4 \pi^2 c^2 v^2$ , in the least squares fit for the  $\text{CF}_3\text{OX}$  series.

Trifluoromethyl hypofluorite ( $\text{CF}_3\text{OF}$ ) is an important reagent in the synthesis of organic and inorganic fluorine compounds (2). There have been some serious spectroscopic studies of this compound since it was first prepared by

Kellog and Cady (11). This compound has been studied by infrared and Raman spectroscopy (4, 12), electron diffraction (13), and microwave spectroscopy (6). Recently Raman spectra have been obtained for  $\text{CF}_3\text{OF}$  in the gas phase as well as diluted in argon matrices at  $8^\circ\text{K}$  (5).

A force constant calculation has been performed by Wilt and Jones based on their assignment and assumed bond lengths and interbond angles (4). This result may not be reliable mainly because of questionable assignments which will be discussed below.

For the OF and CO stretches, Wilt and Jones assigned  $947\text{ cm}^{-1}$  and  $882\text{ cm}^{-1}$ , respectively. After performing their study of  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  in argon matrices at  $8^\circ\text{K}$ , Smardzewski and Fox (5) proposed that these assignments should be reversed. Their new assignments were based on the relative frequencies shift for the CO and OX stretches when going from hypofluorite to hypochlorite. The observation of a shoulder assignable to  $\text{Cl}(37)/\text{Cl}(35)$  isotopic splitting on the band assigned to the OCl stretch in  $\text{CF}_3\text{OCl}$  supported the new assignment.

A torsional frequency of  $56\text{ cm}^{-1}$  was assigned and used to calculate the barrier to internal rotation of the  $\text{CF}_3$  rotor (4). There are some discrepancies between Wilt and Jones' infrared data and ours in the  $300\text{ cm}^{-1}$  to  $200\text{ cm}^{-1}$  region. Since our Raman data are consistent with our IR data, we must question the authenticity of the far infrared data

reported by Wilt and Jones including the  $56\text{ cm}^{-1}$  band.

The potential barrier of  $395\text{ cm}^{-1}$  (1.13 kcal/mole) obtained by Wilt and Jones from the  $56\text{ cm}^{-1}$  band was somewhat lower than the electron diffraction result of  $2.5 \pm 0.5$  kcal/mole (13) and much lower than the microwave result of 3.9 kcal/mole (6) which corresponds to a torsional frequency of approximately  $120\text{ cm}^{-1}$ . A weak band centered at  $126.9\text{ cm}^{-1}$  was observed in the gas phase Raman spectrum of  $\text{CF}_3\text{OF}$  (5). The assignment of this frequency as the  $\text{CF}_3$  torsional mode is the most reasonable choice.

Trifluoromethyl hypochlorite ( $\text{CF}_3\text{OCl}$ ) is the first member of the chloroxy perfluoroalkane series. Molecules known to readily undergo free-radical reactions were very effective in forming derivatives of  $\text{CF}_3\text{OCl}$  in which the  $\text{CF}_3\text{O}$  group is retained (3). The potential of using this compound as a reagent in the syntheses of fluorine compounds is currently being investigated (14).

The preparation, identification and characterization of  $\text{CF}_3\text{OCl}$  were reported almost simultaneously by two separate groups in 1968 and 1969 (3,15). The vibrational assignments were based on infrared spectra only and the main purpose was for characterization. The  $\text{OCl}$  stretching frequency was assigned differently by these two groups. Both  $665\text{ cm}^{-1}$  (15) and  $789\text{ cm}^{-1}$  (3) have been assigned to this mode.

Recently the complementary Raman spectra of  $\text{CF}_3\text{OCl}$

isolated in an argon matrix at 8° K was reported (5). The band at 782.7 cm<sup>-1</sup> was assigned to the OCl stretching mode of CF<sub>3</sub>O<sup>35</sup>Cl on the basis of a low frequency shoulder assignable to the OCl stretching mode of CF<sub>3</sub>O<sup>37</sup>Cl.

No electron diffraction or microwave data for CF<sub>3</sub>OCl have been published to the best of our knowledge. No previous attempt to do a normal coordinate analysis of this compound has been published to the best of our knowledge.

## CHAPTER II EXPERIMENTAL SECTION

### A. Recording of Spectra

#### 1. Materials

Both compounds used in this study were supplied by Dr. D. D. DesMarteau of the Chemistry Department at Kansas State University.  $\text{CF}_3\text{OF}$  was prepared by a similar method to that described in the literature (16). This method involves the cesium fluoride catalyzed addition of fluorine across the carbon-oxygen double bond of  $\text{COF}_2$ .  $\text{CF}_3\text{OCl}$  was prepared using the same method except that  $\text{ClF}$  was used in place of  $\text{F}_2$ .

#### 2. Infrared Instrumentation

Survey and higher resolution spectra from 4000 to 160  $\text{cm}^{-1}$  were obtained with a Perkin-Elmer Model 180 infrared spectrophotometer. Instrument frequency calibration was achieved using known absorption frequencies of polyethylene, teflon, atmospheric water and  $\text{CO}_2$  (17).

Infrared spectra of the vapor were recorded in 10 and 15 cm cells made of Pyrex tubing and a 5 cm cell made of stainless steel. The 15 cm cell with polyethylene windows was used for far-infrared spectra, while the 5 cm and 10 cm cell with  $\text{AgCl}$  windows were used for the mid-range infrared region.

#### 3. Raman Instrumentation

The liquid Raman spectra were recorded in a low tempera-

ture cell similar in design to that of Brown et.al. (18) using a JASCO R-300 Laser Raman Spectrophotometer and the 5145 Å line of a Spectra-Physics Model 164-00 argon ion laser.

The gaseous Raman spectra were recorded at the U. S. Naval Research Laboratory using a Jarrel-Ash Model 500  $\frac{1}{2}$ -meter double monochromator and the 4880 and 5145 Å lines of a Spectra-Physics Model 164-00 argon ion laser (19).

### B. Computer Programs

Several computer programs for the normal coordinate analysis of molecular vibration are available. The ones used in this work are from the set written by Schachtschneider (10).

The programs, CART, GMAT, FPERT and VSEC comprise the package of computer programs selected to fit our needs. CART and GMAT are designed for setting up the vibrational secular equation while VSEC and FPERT are designed for solving the secular equation. FPERT also provides for the least-square refinement of force constants.

CART calculates the Cartesian coordinates of the atoms in a molecule from the bond distances and bond angles. Center of mass and moments of inertia can also be computed. If desired, the principal moments of inertia and the principal Cartesian coordinates are also calculated.

GMAT calculates the Wilson G matrix (i.e., the vibrational inverse kinetic energy matrix) for polyatomic molecules. Input for this program are the Cartesian coordinates (from the output of CART), masses of the atoms, the numbers of the atoms defining a complete set of internal valence coordinates (i.e., bond stretching, valence angle bending, out-of-plane wagging and torsion) (10,20) and the transformation to symmetry coordinates if desired.

VSEC solves the secular equations in internal coordinates by two successive Jacobi diagonalizations. Factoring of the secular equation can be carried out within the program. At the option of the user the mean amplitudes of vibration, the eigenvector inverse matrix  $L^{-1}$ , and Coriolis coefficients may be calculated.

FPERT refines a set of force constants to give a weighted least-squares fit between observed and calculated frequencies. The secular equation in the first cycle is solved in internal coordinates by two successive Jacobi diagonalizations.

All the programs are coded in Fortran IV language. Standardized input-output formats are used so the same cards can be used in all programs.



### CHAPTER III EXPERIMENTAL RESULTS

In this chapter the experimental spectroscopic data for  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  are presented. Summaries derived from both infrared and Raman data are made for both molecules. Only frequencies in the fundamental region ( below  $1400\text{ cm}^{-1}$ ) are included in the summaries.

#### A. Infrared Spectra of $\text{CF}_3\text{OF}$

A summary of the infrared data below  $1400\text{ cm}^{-1}$  is presented in Table 1. The gas-phase infrared spectra of  $\text{CF}_3\text{OF}$  in the fundamental region  $1400\text{-}200\text{ cm}^{-1}$  are shown in Figures 1 and 2. A 10X abscissa scale expansion of the  $429\text{ cm}^{-1}$  band is also shown at the top of Figure 2. A shoulder at  $431\text{ cm}^{-1}$  is observed. This shoulder was first neglected but finally assigned as a fundamental as suggested by the normal coordinate analysis. In general the infrared data presented agree well with that of Wilt and Jones (4). Between  $300$  and  $200\text{ cm}^{-1}$  our spectra differ from those of Wilt and Jones in that our peaks are  $25\text{-}28\text{ cm}^{-1}$  higher in frequencies although the band shapes are somewhat similar.

#### B. Raman Spectra of $\text{CF}_3\text{OF}$

The low and high sensitivity Raman spectra of  $\text{CF}_3\text{OF}$  near  $-196^\circ\text{C}$  are shown in Figures 3 and 4, respectively. A reduced scale Raman gas phase spectrum of  $\text{CF}_3\text{OF}$ , which was provided by Drs. R. R. Smardzewski and W. B. Fox of the



U. S. Naval Research Laboratory, Washington, D. C., is also shown in Figure 3. Experimental conditions including the laser source and temperature are included with each spectrum. Table 1 lists the gas, liquid, and Ar matrix Raman frequencies and the assignments made in this study are also included.

#### C. Infrared Spectra of $\text{CF}_3\text{OCl}$

The infrared spectrum in the fundamental region from  $1400\text{ cm}^{-1}$  to  $180\text{ cm}^{-1}$  in the gas phase is shown in Figures 5 and 6. A summary is in Table 2. The general appearance of the infrared spectrum is very similar to that of Schack and Maya (3). However, the region below  $500\text{ cm}^{-1}$  was not included in their report.

#### D. Raman Spectra of $\text{CF}_3\text{OCl}$

The low and high sensitivity Raman spectra of liquid  $\text{CF}_3\text{OCl}$  at  $-196^\circ\text{C}$  are shown in Figures 7 and 8. The band at  $548\text{ cm}^{-1}$ , which is very prominent on the side of the  $561\text{ cm}^{-1}$  band in the liquid Raman spectrum but completely absent in the IR spectrum, is attributed to residual chlorine complexed with  $\text{CF}_3\text{OCl}$ . Table 2 lists the liquid and Ar matrix Raman frequencies and the assignments made in this study are also included.

#### E. Matrix Spectra of Smardzewski and Fox (5)

Figure 9A illustrates the spectral region containing the  $\nu(\text{OX})$  and  $\nu(\text{CO})$  fundamentals. The complementary matrix infrared spectra are illustrated in the top portions of the

figure. It can readily be seen that the higher frequency band, which is assigned to  $\nu(\text{CO})$ , undergoes a relatively minor shift ( $25.8 \text{ cm}^{-1}$ ) on proceeding from  $\text{CF}_3\text{OF}$  to  $\text{CF}_3\text{OCl}$  while the lower frequency band, which is assigned to  $\nu(\text{OX})$ , experiences a marked decrease in frequency ( $99.9 \text{ cm}^{-1}$ ). Also the chlorine isotopic splitting of the  $782.7 \text{ cm}^{-1}$  matrix Raman band is shown in Figure 9B. The splitting and intensity ratio support the assignment of  $\nu(\text{OCl})$  based on frequency shift. The spectra in Figure 9 are taken from Reference (5).

Table 1. Raman and Infrared Data for  $\text{CF}_3\text{OF}$  below  $1400\text{ cm}^{-1}$ 

Gas		Liquid		$\rho^c$	Ar Matrix	Assignment <sup>d</sup>
Raman <sup>a</sup>	IR <sup>b</sup>	Raman <sup>b</sup>			Raman <sup>a</sup>	
1300	1294	1310			1288	a. $\nu_{\text{CF}_3}(\text{A}')$
1250	1261				1250	a. $\nu_{\text{CF}_3}(\text{A}'')$
1219	1222	1205	0.45		1211	s. $\nu_{\text{CF}_3}$
945	947(Q) <sup>e</sup>	946	0.49		945	$\nu_{\text{CO}}$
881	882(Q) <sup>e</sup>	882	0.03		883	$\nu_{\text{OF}}$
864		868	0.04		871	2 $\delta_{\text{COF}}$
675	678(Q) <sup>e</sup>	679	0.035		678	s. $\delta_{\text{CF}_3}$
606	607	609	0.85		606	a. $\delta_{\text{CF}_3}(\text{A}'')$
581	585	587	0.43		582	a. $\delta_{\text{CF}_3}(\text{A}')$
	431 sh					$\rho_{\text{CF}_3}(\text{A}'')$
429	429(Q) <sup>e</sup>	436	0.32		433	$\delta_{\text{COF}}$
272	278	285??				{ 2 $\tau_{\text{CF}_3}$ and $\rho_{\text{CF}_3}(\text{A}')$ $\tau_{\text{CF}_3}$
246	252	259	0.52		256	
127	-	144	0.86		-	

- a. References (5) and (19).
- b. D. D. DesMarteau and R. M. Hammaker unpublished studies.
- c. These depolarization ratios for the liquids were measured by rotating the polarization of the incident laser beam with no analyzer and no scrambler in the scattered beam. Known depolarized bands below  $500\text{ cm}^{-1}$  gave depolarization ratios between 0.75 and 0.89 with this arrangement. This arrangement is method IV in H. H. Classen, H. Selig, and J. Shamir, *Appl. Spectroscopy*, 23, 8, (1969).
- d. These symbols refer to the work descriptions of the symmetry coordinates defined in Table 4 and of the fundamental vibrations in Table 6.
- e. These are Q branches for bands having PQR structure as follows: 938, 947, 956; 974, 882, 890; 670, 678, 688; 420, 429, 439.

EXPLANATION OF FIGURE 1

Gas-phase Infrared Spectra of  $\text{CF}_3\text{OF}$  in the  
Region  $1400\text{--}500\text{ cm}^{-1}$ .

Spectrometer: Perkin-Elmer Model 180

Sampling Method: 10 cm Path Length Cell with  
AgCl Windows

Pressure: Spectrum A - 3 mm Hg

Spectrum B - 161 mm Hg

Energy Mode: Constant  $I_0$

Resolution: Spectrum A -  $1.1\text{ cm}^{-1}$  at  $1400\text{ cm}^{-1}$   
Spectrum B -  $1.0\text{ cm}^{-1}$  at  $1000\text{ cm}^{-1}$

Gain: 6


Slit Program: 5

Time Constant: 1

Scan Time: Fine 5, Coarse 10

Suppression: 5

Temperature:  $25^\circ\text{ C}$



EXPLANATION OF FIGURE 2

Gas-phase Infrared Spectra of  $\text{CF}_3\text{OF}$  in the  
Region  $650\text{-}200\text{ cm}^{-1}$ .

Spectrometer: Perkin-Elmer Model 180

Sampling Method: 15 cm Path Length Cell with  
Polyethylene Windows

Pressure: 438 mm Hg

Energy Mode: Constant  $I_0$

Resolution:  $3.0\text{ cm}^{-1}$  at  $650\text{ cm}^{-1}$

Gain: 6

Slit Program: 5

Time Constant: 1

Scan Time: Fine 5, Coarse 10

Suppression: 5

Temperature:  $25^\circ\text{C}$

A 10X abscissa scale expansion of the  $429\text{ cm}^{-1}$   
band is shown directly above. A shoulder at  $431\text{ cm}^{-1}$   
is observed.

### EXPLANATION OF FIGURE 3

A. Raman Spectrum of Liquid  $\text{CF}_3\text{OF}$

Spectrometer: JASCO Model R-300

Slit Setting:  $150\ \mu\text{m}$

Sensitivity: 1

Response: 0.5 Seconds

Scan Rate: 4

Power: 140 mW

Source:  $\text{Ar}^+$  Laser at  $5145\ \text{\AA}$

Temperature:  $-196^\circ\text{C}$

B. Raman Spectrum of Gaseous  $\text{CF}_3\text{OF}$

(Courtesy of Dr. Richard R. Smardzewski of  
Naval Research Laboratory (19))

Spectrometer: Jarrel-Ash Model 500

Slit Setting:  $100\ \mu\text{m}$

Concentration: 150 mm Hg

Sensitivity: 800


Time Constant: 1 Second

Scan Rate: 2.5 cm/sec

Source:  $\text{Ar}^+$  Laser at  $5145\ \text{\AA}$

Power: 1.5 W

Temperature:  $25^\circ\text{C}$





EXPLANATION OF FIGURE 4

High Sensitivity Raman Spectra of Liquid  $\text{CF}_3\text{OF}$

Spectrometer: JASCO Model R-300

Slit Setting:  $150\ \mu\text{m}$

Sensitivity: 5

Response: 2 Seconds

Scan Rate: 4

Source:  $\text{Ar}^+$  Laser at  $5145\ \text{\AA}$

Power: 140 mW

Temperature:  $-196^\circ\text{C}$

EXPLANATION OF FIGURE 5

Gas-phase Infrared Spectra of  $\text{CF}_3\text{OCl}$  in the  
Region  $1400\text{--}500\text{ cm}^{-1}$ .

Spectrometer: Perkin-Elmer Model 180

Sampling Method: 10 cm Path Length Cell with  
AgCl Windows

Pressure: Spectrum A - 3 mm Hg

Spectrum B - 65 mm Hg

Energy Mode: Constant  $I_0$

Resolution: Spectrum A -  $1.7\text{ cm}^{-1}$  at  $1300\text{ cm}^{-1}$

Spectrum B - Slit width 0.2 mm at  
 $4000\text{ cm}^{-1}$

Gain: 6


Slit Program: 5

Time Constant: 1

Scan Time: Fine 5, Coarse 10

Suppression: 5

Temperature:  $25^\circ\text{C}$



EXPLANATION OF FIGURE 6

Gas-phase Infrared Spectra of  $\text{CF}_3\text{OCl}$  in the  
Region  $650\text{-}180\text{ cm}^{-1}$ .

Spectrometer: Perkin-Elmer Model 180

Sampling Method: 15 cm Path Length Cell with  
Polyethylene Windows

Pressure: Spectrum A - 100 mm Hg

Spectrum B - 140 mm Hg

Energy Mode: Constant  $I_0$

Resolution: Spectrum A -  $3.4\text{ cm}^{-1}$  at  $650\text{ cm}^{-1}$

Spectrum B -  $2.0\text{ cm}^{-1}$  at  $300\text{ cm}^{-1}$


Gain: 6

Slit Program: 5

Scan Time: Fine 5, Coarse 10

Suppression: 5

Temperature:  $25^\circ\text{C}$



EXPLANATION OF FIGURE 7

Raman Spectrum of Liquid  $\text{CF}_3\text{OCl}$

Spectrometer: JASCO Model R-300

Slit Setting:  $150\ \mu\text{m}$

Sensitivity: 3


Response: 0.5 Seconds

Scan Rate:  $\times 4$

Source:  $\text{Ar}^+$  Laser at  $5145\ \text{\AA}$

Power: 120 mW

Temperature:  $-196^\circ\text{C}$



EXPLANATION OF FIGURE 8

High Sensitivity Raman Spectra of Liquid  $\text{CF}_3\text{OCl}$

Spectrometer: JASCO Model R-300

Slit Setting:  $150\ \mu\text{m}$

Sensitivity: 5

Response: 2 Seconds

Scan Rate: 4

Source:  $\text{Ar}^+$  Laser at  $5145\ \text{\AA}$

Power: 240 mW

Temperature:  $-196^\circ\text{C}$



$\Delta V \text{ (CM}^{-1}\text{)}$


200

400

600

800


INTENSITY



EXPLANATION OF FIGURE 9

$\nu(\text{OX})$  and  $\nu(\text{CO})$  Stretches for  $\text{CF}_3\text{OX}$  Molecules.  
(Taken from Reference 5)

- A. Figure A. Argon matrix Raman and infrared spectra of  $\text{CF}_3\text{OX}$  ( $\text{X}=\text{F}, \text{Cl}$ ) in the  $\nu(\text{OX})$  and  $\nu(\text{CO})$  fundamental region. (a)  $\text{Ar}/\text{CF}_3\text{OF}=100$ , (b)  $\text{Ar}/\text{CF}_3\text{OCl}=100$ . Raman spectra were run with  $5145 \text{ \AA}$  excitation (1.7 W), 400 CPS amplification,  $5 \text{ cm}^{-1}$  resolution, 0.25 cm/sec scan speed and 5 sec time constant.
- B. Figure B. Chlorine isotopic splitting of the  $782.7 \text{ cm}^{-1}$  Raman band of  $\text{CF}_3\text{OCl}$  in an argon matrix at  $8^\circ\text{K}$ .  $\text{Ar}/\text{CF}_3\text{OCl}=100$ ,  $5145 \text{ \AA}$  excitation (1.7 W), 100 CPS amplification,  $4 \text{ cm}^{-1}$  resolution, 0.05 cm/sec scan speed, 10 sec time constant.



#### CHAPTER IV NORMAL COORDINATE ANALYSIS

Having completed the vibrational assignment for the  $\text{CF}_3\text{OX}$  series, it is now possible to carry out a normal coordinate analysis. The purpose of this calculation is to see whether or not the vibrational frequencies chosen as fundamentals in the new assignments can be fitted reasonably well with a realistic potential function. If so, this may be interpreted as evidence for the correctness of the assignment. A second purpose is to clarify the descriptions of the motion in each vibrational mode. A large amount of mixing and very complicated normal coordinates are expected due to the closeness of the masses of atoms in these two molecules.

##### A. Structural Models

$\text{CF}_3\text{OF}$  has been shown in a recent electron diffraction study to possess  $\text{C}_s$  symmetry (13). The geometric parameters of  $\text{CF}_3\text{OF}$  used in this work are listed in Table 3.

Unfortunately no electron diffraction or microwave data for  $\text{CF}_3\text{OCl}$  have been published. Thus, it is necessary to assume bond lengths and interbond angles.  $\text{CF}_3\text{OCl}$  is assumed to be of  $\text{C}_s$  symmetry with the  $\text{CF}_3\text{O}$  portion of the same geometry as that of  $\text{CF}_3\text{OF}$ . The O-Cl bond related geometry for  $\text{CF}_3\text{OCl}$  is deduced by comparison of  $\text{CF}_3\text{OF}$  with  $\text{F}_2\text{O}$  and  $\text{F}_2\text{O}$  with  $\text{Cl}_2\text{O}$ . Since the O-F bonds in  $\text{CF}_3\text{OF}$  and  $\text{F}_2\text{O}$  differ by only about

0.01 Å, the O-Cl bond length from  $\text{Cl}_2\text{O}$  is used for  $\text{CF}_3\text{OCl}$ . Since the Cl-O-Cl angle in  $\text{Cl}_2\text{O}$  is  $8^\circ$  larger than the F-O-F angle in  $\text{F}_2\text{O}$ , the C-O-Cl angle in  $\text{CF}_3\text{OCl}$  is taken as  $8^\circ$  larger than the C-O-F angle in  $\text{CF}_3\text{OF}$ . The geometric parameters of  $\text{CF}_3\text{OCl}$  are also shown in Table 3, and the structure of  $\text{CF}_3\text{OX}$  is shown in Figure 10.

#### B.. Construction of the G Matrix

In order to construct the G matrices, it is necessary to know the structures of the molecules, the atomic masses, and a set of internal coordinates.

The G matrices used in this study were computed using the programs CART and GMAT which were described in Chapter II. Geometry used for  $\text{CF}_3\text{OF}$  was determined from electron diffraction. Estimated geometry was used for  $\text{CF}_3\text{OCl}$ . These structural parameters are all listed in Table 3 in the previous section. The atomic masses were taken from the carbon 12 system:

$$\begin{aligned}\text{C}(12) &= 12.00000 \\ \text{F}(19) &= 18.99840 \\ \text{O}(16) &= 15.99491 \\ \text{Cl}(35) &= 34.96885\end{aligned}$$

The G matrix elements were not weighted. Symmetrizations were accomplished using symmetry coordinates shown in Table 4. The internal coordinates used to generate these symmetry coordinates are shown in Figure 11.

Table 3. Structural Data for  $\text{CF}_3\text{OX}$  Compounds

Geometrical Parameters <sup>a</sup>	$\text{CF}_3\text{OF}^b$	$\text{CF}_3\text{OCl}^c$
$\text{R}(\text{C}-\text{F})(\text{\AA})$	1.319	1.319
$\text{R}(\text{C}-\text{O})(\text{\AA})$	1.395	1.395
$\text{R}(\text{O}-\text{X})(\text{\AA})$	1.421	1.700
$\alpha(\text{deg})$	109.4	109.4
$\beta(\text{deg})$	4.1	4.1
$\gamma(\text{deg})$	104.8	112.8
$\delta(\text{deg})$	70.5	70.5
$\theta(\text{deg})$	105.4	105.4
$I_a(\text{amu}-\text{\AA}^2)$	89.6	89.6
$I_b(\text{amu}-\text{\AA}^2)$	164.6	254.9
$I_c(\text{amu}-\text{\AA}^2)$	166.0	256.4
$\kappa$	-0.98	-0.99


- a. The angles  $\alpha, \beta, \gamma, \delta$ , and  $\theta$  are illustrated in Figure 10. The asymmetry parameter,  $\kappa$ , is defined as  $\kappa = (2B-A-C)/(A-C)$  where A, B, C are rotational constants proportional to  $(I_a)^{-1}, (I_b)^{-1}, (I_c)^{-1}$ , respectively. A  $\kappa$  value of -1 corresponds to a prolate symmetric top.
- b. The geometry for  $\text{CF}_3\text{OF}$  was determined from the electron diffraction data of<sup>3</sup>Ref. 13.
- c. The geometry for  $\text{CF}_3\text{OCl}$  was estimated as discussed in the text. The data used<sup>3</sup>for  $\text{Cl}_2\text{O}$  and  $\text{F}_2\text{O}$  were from L. E. Sutton, ed. "Tables of Interatomic<sup>2</sup>Distances and Configuration in Molecules and Ions", Special Publication No. 11, The Chemical Society, London (1958) p. M67.

# EXPLANATION OF FIGURE 10

## Structure of $\text{CF}_3\text{OX}$ Molecules.

Structural data for both molecules are listed in Table 3.

The dashed line (-----) indicates the  $\text{C}_3$  axis of the  $\text{CF}_3$  group which has local  $\text{C}_{3v}$  symmetry. The angle of tilt,  $\beta$ , is in the FCOX plane and is the angle between the  $\text{C}_3$  axis and the C-O bond. For a positive angle of tilt like the one shown in this figure the F atom in the FCOX plane is closer to the O atom than are the two out of plane F atoms.





### C. Utilization of Symmetry

Prior to solving the secular equation, the symmetry of the molecular system may be introduced to advantage as a means of simplifying the secular determinant. Symmetry coordinates, which can be constructed from a suitable linear combination of internal coordinates, are used to factor the secular determinant into smaller, square subblocks. The vibrational problem then reduces to the solution of a series of independent, smaller secular determinants.

For the assumed  $C_s$  symmetry of the  $CF_3OX$  series, group theory analysis (21) predicts twelve normal modes of vibration, eight of species  $A'$  (symmetric with respect to the plane of symmetry) and four of species  $A''$  (antisymmetric with respect to the plane of symmetry). All twelve are infrared and Raman active, with the  $A'$  modes being polarized and  $A''$  modes depolarized in the Raman spectrum (4). The symmetry coordinates used to factor the  $F$  and  $G$  matrices are given in Table 4 in terms of the internal coordinates shown in Figure 11.

### D. Choice of Force Fields

The number of vibrational frequencies for a single polyatomic molecule is invariably less than the number of force constants in the general harmonic force field. It was obviously impossible to determine any more than a very limited force field for the  $CF_3OX$  series because of the

paucity of data. Since certain interaction constants cannot be determined with any degree of certainty even with a large amount of data (22), we can assume a simplified force field such as the valence force field with selected interaction constants set equal to zero.

The F matrix used in this study was set up in terms of internal coordinates and listed in array form in Figure 12. Symmetrization was obtained by using the symmetry coordinates listed in Table 4. The least squares adjustment of force constants were carried out using the program FPRT described earlier.

Results of the force constant analysis for  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  are shown in Tables 5, 6, and 7. The meaning of 'Chemical Fitting' and 'Mathematical Fitting' appearing in Tables 6 and 7 will be explained in the next section.

#### E. Chemical Approach vs. Mathematical Approach

A complete and reliable specification of the potential energy of a molecular system serves as the primary goal of a normal coordinate analysis. The most common approach toward evaluating a potential function from observed vibrational frequencies involves some method of force constant adjustment such that the calculation reproduces the observed data as closely as possible (22-26). In most cases, harmonic frequencies are not available and consequently anharmonicity effects are an additional source of error. We are using

experimental frequencies not harmonic frequencies in this work.

The method of least-squares fitting, which was employed in this study, represents a particularly effective procedure for refining an initial set of force constants to optimize the fit between the calculated and observed frequencies. This mathematical approach, as we may put it, regards the vibrational analysis as a problem in applied mathematics and there is no relationship between force constant values and chemistry. Since the method for adjusting force constants does not insure a unique potential function, other molecular information may be used as constraints in limiting the acceptable solutions of the secular equation(1).

The imposed constraints represent a kind of chemical approach. In this approach the chemistry of this molecule should be taken into consideration. The force constants must be consistent with other data about the compound, including the known chemical properties of this compound and with a related series of compounds it should be possible to transfer force constants for similar group. Although the transferability of force constants is not always expected to be strictly observed, it should be consistent within certain ranges obtained from similar molecules.

Without proper choice and restraint on the major diagonal force constants, the mathematical fitting minimum found by

least-squares analysis may be just another incorrect minimum on the fit vs. force constant set surface. The fitting may be greatly improved with no restriction on major diagonal force constants. The final mathematical fitting solution certainly need not correspond to the real molecule and this solution may not have any chemical significance.

From our work the difference of these two approaches is very clear in the case of  $\text{CF}_3\text{OCl}$ . The mathematical fitting is excellent, but the force constant for OCl stretching must be raised to an unreasonably high value to get this excellent fit. All the discussion in the next chapter is based entirely on the results of chemical fitting.

Table 4. Symmetry Coordinates for  $\text{CF}_3\text{OX}$  Molecules  
of  $\text{C}_s$  Symmetry<sup>a</sup>


Antisymmetric $\text{CF}_3$ stretch	$S_1 = \sqrt{6}^{-1} (2\Delta r_4 - \Delta r_5 - \Delta r_6)$
Symmetric $\text{CF}_3$ stretch	$S_2 = \sqrt{3}^{-1} (\Delta r_4 + \Delta r_5 + \Delta r_6)$
OX stretch	$S_3 = \Delta d$
Antisymmetric $\text{CF}_3$ deformation	$S_4 = \sqrt{6}^{-1} (2\Delta \alpha_4 - \Delta \alpha_5 - \Delta \alpha_6)$
A' Symmetric $\text{CF}_3$ deformation	$S_5 = \sqrt{6}^{-1} (\Delta \alpha_4 + \Delta \alpha_5 + \Delta \alpha_6 - \Delta \beta_4 - \Delta \beta_5 - \Delta \beta_6)$
COX bend	$S_6 = \Delta \gamma$
CO stretch	$S_7 = \Delta l$
$\text{CF}_3$ rock (in-plane)	$S_8 = \sqrt{6}^{-1} (2\Delta \beta_4 - \Delta \beta_5 - \Delta \beta_6)$
Redundant	$S_9^b = \sqrt{6}^{-1} (\Delta \alpha_4 + \Delta \alpha_5 + \Delta \alpha_6 + \Delta \beta_4 + \Delta \beta_5 + \Delta \beta_6)$
Antisymmetric $\text{CF}_3$ stretch	$S_{10} = \sqrt{2}^{-1} (\Delta r_5 - \Delta r_6)$
Antisymmetric $\text{CF}_3$ deformation	$S_{11} = \sqrt{2}^{-1} (\Delta \alpha_5 - \Delta \alpha_6)$
A'' $\text{CF}_3$ rock (out-of-plane)	$S_{12} = \sqrt{2}^{-1} (\Delta \beta_5 - \Delta \beta_6)$
$\text{CF}_3$ torsion	$S_{13} = \Delta \tau$

- a. There will be three related quantities having numbering systems as follows: these thirteen symmetry coordinates,  $S_i$ , (including one redundancy in the A' symmetry block), twelve coresponding normal coordinates,  $Q_i$ , in Tables 10 and 11, and twelve normal mode frequencies,  $\nu_i$ , in Tables 6, 7, 9, 12, 13, and 14. The relation between the three numbering systems is given in Table 6.
- b. These coordinates are correct for tetrahedral  $\text{C}_{3v}$  angles. However, they may be used for the non-tetrahedral angle case since the redundancy is removed during diagonalization of the G matrix.

EXPLANATION OF FIGURE 11

Internal Coordinates for  $\text{CF}_3\text{OX}$  Molecules.

$\text{C}_s$  symmetry is assumed with the F, C, O,  
and X atoms in the same plane.



# EXPLANATION OF FIGURE 12

Unsymmetrized F Matrix for  $\text{CF}_3\text{OX}$  Molecules.

Internal coordinates are from Figure 11. The F matrix elements  $r\alpha$  and  $\alpha\beta$  are for three opposite pairs of internal coordinates. The F matrix elements  $r\alpha'$  and  $\alpha\beta'$  are for six adjacent pairs of internal coordinates. The F matrix elements  $r\beta$  are for three adjacent pairs of internal coordinates. The F matrix elements  $r\beta'$  are for six opposite pairs of internal coordinates.



[illegible]

Table 5. Internal Coordinate Force Constants<sup>a, b</sup> for CF<sub>3</sub>OF and CF<sub>3</sub>OCl

Internal Force Constant <sup>c</sup>	CF <sub>3</sub> OF		CF <sub>3</sub> OdL		CF <sub>3</sub> OCl	
	Chemical Fitting	Mathematical Fitting	Chemical Fitting	Mathematical Fitting	Chemical Fitting	Mathematical Fitting
r	6.950	6.775	6.950		6.846	
l	5.200	5.656	5.200		5.877	
d	3.850±0.220	3.785±0.035	2.800±0.556		3.885±0.063	
α	2.000	1.980	2.050		1.836	
β	1.200	1.235	1.150		1.266	
γ	1.800±0.172	1.918±0.027	1.700±0.377		2.014±0.042	
τ	0.0167±0.0001	0.0165±0.0001	0.0161±0.0019		0.0165±0.0002	
r r	0.900	1.120	0.900		0.734	
r l	0.850	0.850	0.900		0.850	
l d	1.000±0.243	0.425±0.095	0.300±1.294		0.896±0.199	
r α (opp.)	0.120	0.120	0.120		0.120	
r α' (adj.)	0.910	0.950	0.910		1.040	
r β (adj.)	0.750	0.500	0.700		0.775	
r β' (opp.)	0.100	0.100	0.100		0.100	
l α	0.200	0.200	0.200		0.200	
l β	0.600	0.600	0.600		0.600	
l γ	0.000±0.245	0.614±0.055	0.100±0.933		0.816±0.113	
d γ	0.250±0.200	0.456±0.027	0.100±0.333		0.918±0.041	

34a

Internal Force Constant <sup>c</sup>	CF <sub>3</sub> OF		CF <sub>3</sub> OF		CF <sub>3</sub> OC1		CF <sub>3</sub> OC1	
	Chemical	Fitting	Mathematical	Fitting	Chemical	Fitting	Mathematical	Fitting
$\alpha \alpha$	0.300		0.300		0.350		0.300	
$\alpha \beta$ (opp.)	0.000		0.000		0.000		0.000	
$\alpha \beta'$ (adj.)	0.150		0.150		0.150		0.150	
$\beta \beta$	0.100		0.100		0.100		0.100	

- a. Stretching constants in units of millidynes per angstrom. Stretch-bend interaction constants in units of millidynes per radian. Bending and torsion constants in units of millidyne angstrom per square radian.
- b. The force constants  $d$ ,  $\gamma$ ,  $\tau$ ,  $ld$ ,  $l\gamma$ , and  $d\gamma$  associated with the OX part of the CF<sub>3</sub>OX structure are likely to change from CF<sub>3</sub>OF to CF<sub>3</sub>OC1 and are of primary concern. The remaining force constants have been transferred or estimated using CF<sub>3</sub>H (53) and CF<sub>3</sub>H (54). Error estimates corresponding to a set of force constants can be obtained for a restricted number of members of the set by using the set as input to the FPRT program with the restricted number of members of the set allowed to vary and with the number of perturbations restricted to zero. The errors shown for  $d$ ,  $\gamma$ ,  $\tau$ ,  $ld$ ,  $l\gamma$  and  $d\gamma$  are obtained this way and are the errors listed as \* in the FPRT output and calculated using  $\lambda_i(\text{obs}) - \lambda_i(\text{calc})$  where  $\lambda = 4\pi^2 c^2 \nu^2$  (see numbered statement 62 in the FINI subroutine of FPRT in Tech. Report No. 57-65 in. reference 10).
- c. See Figures 11 and 12 for definition of these internal coordinates.

Table 6. Assignments of the Fundamental Vibrations<sup>a</sup>  
of Trifluoromethyl hypofluorite ( $\text{CF}_3\text{OF}$ )

$\nu_i$	Approximate description	Obs <sup>b</sup>	Chemical Fitting	Mathematical Fitting
A' Species				
1	Antisym $\text{CF}_3$ stretch	1294	1321	1299
2	Sym $\text{CF}_3$ stretch	1222	1214	1222
3	CO stretch	947	941	948
4	OF stretch	882	883	882
5	Sym $\text{CF}_3$ deformation	678	674	678
6	Antisym $\text{CF}_3$ deformation	585	589	585
7	COF bend	429	420	429
8	In-plane $\text{CF}_3$ rock	278	267	278
A'' Species				
9	Antisym $\text{CF}_3$ stretch	1261	1256	1251
10	Antisym $\text{CF}_3$ deformation	607	616	608
11	Out-of-plane $\text{CF}_3$ rock	431	429	431
12	$\text{CF}_3$ torsion	127	128	127

Average error

1.15%

0.137%

- a. The numbering system for the normal mode frequencies,  $\nu_i$ , normal coordinates,  $Q_i$ , and symmetry coordinate,  $S_i$ , are related as follows:


$\nu_i$  1 2 3 4 5 6 7 8 - 9 10 11 12

$Q_i$  1 2 3 4 5 6 7 8 - 9 10 11 12

$S_i$  1 2 7 3 5 4 6 8 9 10 11 12 13

As shown in Table 4,  $S_9$  is the redundant symmetry coordinate in the A' symmetry block.

- b. Frequencies are from gas phase infrared spectra (Table 1) except the  $\text{CF}_3$  torsion is from gas phase Raman spectra (5, 19).






Table 2. Raman and Infrared Data for  $\text{CF}_3\text{OCl}$  below  $1400\text{ cm}^{-1}$ 

Gas IR <sup>a</sup>	Liquid Raman <sup>a</sup>	$\rho$ <sup>b</sup>	Ar Matrix Raman <sup>c</sup>	Assignments <sup>d</sup>
1271	1275	0.60	1269	$\text{a.v}_{\text{CF}_3}(\text{A}')$
1230			1221	$\text{a.v}_{\text{CF}_3}(\text{A}'')$
1213	1190	0.66	1200	$\text{s.v}_{\text{CF}_3}$
919(Q) <sup>e</sup>	917	0.46	920	$\text{v}_{\text{CO}}$
780	781	0.12	783	$\text{v}_{\text{OCl}}^{35}$
			776	$\text{v}_{\text{OCl}}^{37}$
665(Q) <sup>e</sup>	666	0.38	663	$\text{s.}\delta_{\text{CF}_3}$
609	611	0.76	609	$\text{a.}\delta_{\text{CF}_3}(\text{A}'')$
557	561	0.26	558	$\text{a.}\delta_{\text{CF}_3}(\text{A}')$
	430	?		$\rho_{\text{CF}_3}(\text{A}'')$
393(Q) <sup>e</sup>	397	0.27	398	$\delta_{\text{COCl}}$
233	233?		239	{ $2\tau_{\text{CF}_3}$ and $\rho_{\text{CF}_3}(\text{A}')$ $\tau_{\text{CF}_3}$
220??	220	0.47		
-	122	0.81		

- a. D. D. DesMarteau and R. M. Hammaker, unpublished studies.
- b. These depolarization ratios for the liquids were measured by rotating the polarization of the incident laser beam with no analyzer and no scrambler in the scattered beam. Known depolarized bands below  $500\text{ cm}^{-1}$  gave depolarization ratios between 0.75 and 0.89 with this arrangement. The arrangement is method IV in H. H. Claassen, H. Selig, and J. Shamir, *Appl. Spectroscopy*, **23**, 8 (1969).
- c. References (5) and (19).
- d. These symbols refer to the word description of the symmetry coordinates defined in Table 4 and of the fundamental vibrations in Table 6.
- e. These are Q branches for bands having PQR structure as follows: 915, 919, 923; 659, 665, 671; 386, 393, 399.

Table 7. Assignments of the Fundamental Vibrations<sup>a</sup>  
of Trifluoromethyl Hypochlorite ( $\text{CF}_3\text{OCl}$ )

$\nu_i$	Approximate Description	Obs <sup>b</sup>	Chemical Fitting	Mathematical Fitting
A' Species				
1	Antisym $\text{CF}_3$ stretch	1271	1309	1273
2	Sym $\text{CF}_3$ stretch	1213	1208	1213
3	CO stretch	919	935	919
4	OCl stretch	780	774	783
5	Sym $\text{CF}_3$ deformation	665	658	664
6	Antisym $\text{CF}_3$ deformation	557	571	558
7	COCl bend	393	348	392
8	In-plane $\text{CF}_3$ rock	233	217	233
A'' Species				
9	Antisym $\text{CF}_3$ stretch	1230	1262	1229
10	Antisym $\text{CF}_3$ deformation	609	613	608
11	Out-of-plane $\text{CF}_3$ rock	430	422	430
12	$\text{CF}_3$ torsion	108	108	110
Average error			2.72%	0.265%

a. See footnote a in Table 6.

b. Frequencies are from gas phase infrared spectra (Table 2) with the following exceptions:  $430\text{ cm}^{-1}$  from liquid Raman and  $108\text{ cm}^{-1}$  estimated as liquid Raman less 11.8% ( $122\text{ cm}^{-1} - 14\text{ cm}^{-1} = 108\text{ cm}^{-1}$ ) since  $\text{CF}_3\text{OF}$  has gas phase Raman frequency 11.8% below liquid Raman frequency for the  $\text{CF}_3$  torsion. See footnotes a and b in Table 9.



CHAPTER V  
RESULTS, CONCLUSION, AND OTHER REMARKS

Results of the force constant analyses of  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  are shown in Tables 5, 6 and 7.

Numerical values of the force constants are reproduced in Table 5 whereas the calculated fundamental frequencies are compared with the observed values in Tables 6 and 7.

A. Comments on the Choice of Force Fields

It can be seen that the agreement of calculated and observed frequencies for the twelve fundamental modes is 'good' for chemical fitting but 'excellent' for mathematical fitting. If the correctness of force field is measured by the accuracy of the frequency fit, it is very obvious what the choice should be.

But for a case like ours where  $\text{CF}_3\text{OF}$ ,  $\text{CF}_3\text{OCl}$ ,  $\text{CF}_3\text{OOH(D)}$ ,  $\text{CF}_3\text{OOF}$ , and  $\text{CF}_3\text{OOC}$ l all have a  $\text{CF}_3\text{O}$  fragment in common, the explicit recognition of the importance of chemical factors become increasingly pertinent to normal coordinate analysis and will be the criterion of a reasonable force field.

$\text{CF}_3\text{OF}$  is thermally stable in the IR cell at room temperature and the gas phase Raman spectra can be obtained at room temperature with  $\text{Ar}^+$  laser radiation.  $\text{CF}_3\text{OCl}$  is stable at room temperature only under scrupulously dry, inert condition. Photolytic decomposition of  $\text{CF}_3\text{OCl}$  is quite rapid and the decomposition products found are consistent with a

two-step reaction path involving the two radicals  $\text{CF}_3\cdot$  and  $\text{Cl}\cdot$  (3). In the course of this work, the  $\text{CF}_3\text{OCl}$  began to decompose during the recording of its IR spectrum at ambient temperature. Smardzewski and Fox (5) reported that  $\text{CF}_3\text{OCl}$  photolyzed almost immediately upon trying to record its gas phase Raman spectrum at room temperature with an  $\text{Ar}^+$  laser.

This evidence suggests that the O-F bond in  $\text{CF}_3\text{OF}$  may be stronger than the O-Cl bond in  $\text{CF}_3\text{OCl}$ . Chemical consideration would prohibit the use of about the same values for the force constants of the O-X bond. Then identical or very similar values for the force constants related to the  $\text{CF}_3\text{O}$  fragment should be observed while the O-X bond related force constants should be allowed to vary in  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  independently to improve the fit in both molecules.

Pierce, Di Cianni, and Jackson (27) as well as Morino and Saito (28) have done a good analysis on  $\text{F}_2\text{O}$  and the OF stretching force constant  $d(\text{OF})$  was determined to be 3.78 or 3.95 or 4.10 mdyne/Å by several methods.

The  $d(\text{OF})$  values in  $\text{HOF}$  obtained by Noble and Pimentel (29) and Ogilvie (30) are 4.37 and 4.27 mdyne/Å, respectively. Noble and Pimentel have suggested that  $d(\text{OF})$  is larger in  $\text{HOF}$  than in  $\text{F}_2\text{O}$  because the electronegativity of fluorine weakens the O-H bond and strengthens the O-F bond by attraction of negative charge out of the O-H bond into the O-F bond. For such a charge competition in  $\text{CF}_3\text{OF}$ , it seems reasonable that

the  $\text{CF}_3$  group would be better able to compete with the F atom for charge over the  $\text{CF}_3\text{OF}$  frame. Thus  $d(\text{OF})$  in  $\text{CF}_3\text{OF}$  should be smaller than that in  $\text{HOF}$ . The value of  $3.85 \text{ mdyn/\AA}$  used in this study can be considered to be a chemically reasonable value, since it is in the range for  $\text{F}_2\text{O}$ .

Rochkind and Pimentel(31) should have the most reliable force field for  $\text{Cl}_2\text{O}$  in view of their isotopic data. The low value of  $2.75 \text{ mdyn/\AA}$  for  $d(\text{OCl})$  in  $\text{Cl}_2\text{O}$  is commensurate with the relatively long Cl-O bond of  $1.70 \text{ \AA}$ . This value was proclaimed to be the prototype force constant for a normal Cl-O single bond (31). Repetition of the calculation on  $\text{Cl}_2\text{O}$  gave  $d(\text{OCl})$  the values of  $2.75(32)$  and  $2.88(33) \text{ mdyn/\AA}$  respectively. A value of  $2.65 \text{ mdyn/\AA}$  was obtained for  $d(\text{OCl})$  in  $\text{ClO}_3\text{OCl}(34)$ . The work of Witt and Hammaker(35) on the skeletal mode of  $\text{Cl}_2\text{O}_7$  put a value of  $3.09 \text{ mdyn/\AA}$  on the skeletal OCl stretch.

Bond-stretching force constants,  $d(\text{OF})$  and  $d(\text{OCl})$ , taken from related molecules containing O-F and O-Cl bonds are listed in Table 8 for comparison.

The electronegativity arguments of Noble and Pimentel can be applied to the  $d(\text{OCl})$  value in  $\text{HOCl}$  and  $\text{CF}_3\text{OCl}$  also. Certainly, the chlorine is electronegative enough relative to hydrogen to attract charge out of the O-H bond into the Cl-O bond. Thus  $d(\text{OCl})$  in  $\text{HOCl}$  should be larger than in  $\text{Cl}_2\text{O}$ . One might argue that replacement of H by  $\text{CF}_3$  would shift

some charge away from the Cl-O bond into the C-O bond as the  $\text{CF}_3$  group would be better able to compete with Cl for charge over the C-O-Cl frame than would H over the HOCl frame. Thus  $d(\text{OCl})$  in  $\text{CF}_3\text{OCl}$  should be smaller than in HOCl.

From Table 8 we can observe that the previous work is generally consistent with  $d(\text{OF})$  larger than  $d(\text{OCl})$  with the difference ranging from 0.6 to 1.2  $\text{mdyn}/\text{\AA}$ . Judging from the above argument, we believe the values of  $d(\text{OX})$  of  $\text{CF}_3\text{OX}$  should close to those of  $\text{X}_2\text{O}$ . The force field with  $d(\text{OCl})$  higher than or close to  $d(\text{OF})$  should be dismissed.

Table 8. Bond-stretching Force Constants of O-F and O-Cl Bonds from related Molecules

Compound	Force Constants	
	$d(\text{OF})(\text{mdyn}/\text{\AA})$	$d(\text{OCl})(\text{mdyn}/\text{\AA})$
$\text{F}_2\text{O}^{\text{a}}$	3.95	
$\text{Cl}_2\text{O}^{\text{b}}$		2.75
$\text{ClO}_3\text{OF}^{\text{c}}$	3.56	
$\text{ClO}_3\text{OCl}^{\text{c}}$		2.65
$\text{HOF}^{\text{d}}$	4.27	
$\text{HOCl}^{\text{d}}$		3.68

a. Data from Ref. 27.

b. Data from Ref. 31.

c. Data from Ref. 35.

d. Data from Ref. 30.

## B. Comments on Assignments Implied by Force Constant Calculation

Three fundamentals are expected in the region between 1180 and 1300  $\text{cm}^{-1}$  for the  $\text{CF}_3$  stretching modes ( $\nu_{\text{CF}_3}$ ) of the  $\text{CF}_3\text{OX}$  series. Two of species A' and one of species A'' are expected under  $C_s$  symmetry. Our calculations gave a better fit for  $\nu_{\text{CF}_3}$  if the highest and lowest frequencies are in the A' block and middle frequency is in the A'' block. This indicates that the assignments of the middle frequency (1261  $\text{cm}^{-1}$ ) in the A' block and lowest frequency (1222  $\text{cm}^{-1}$ ) in the A'' block for antisymmetric  $\text{CF}_3$  stretching by Wilt and Jones (4) should be reversed. This sequence is used in final assignments of six compounds in both the  $\text{CF}_3\text{OX}$  and  $\text{CF}_3\text{OOX}$  series (8).

Initially our sample of  $\text{CF}_3\text{OF}$  showed many Raman bands in addition to those in Figures 3 and 4. These were shown to be due to  $\text{CF}_2(\text{OF})_2$ . In the preparation of  $\text{CF}_3\text{OF}$  by addition of  $\text{F}_2$  to  $\text{COF}_2$ ,  $\text{CO}_2$  impurity in the  $\text{COF}_2$  can add two moles of  $\text{F}_2$  to form  $\text{CF}_2(\text{OF})_2$ . The sample of  $\text{CF}_3\text{OF}$  in Figures 3 and 4 has had most of the  $\text{CF}_2(\text{OF})_2$  separated by low temperature chromatography and only two small peaks remain corresponding to strong bands from  $\text{CF}_2(\text{OF})_2$ . Following this purification, a band at 868  $\text{cm}^{-1}$  remained which was not due to  $\text{CF}_2(\text{OF})_2$ . Smardzewski and Fox (5,19) report an analogous band for  $\text{CF}_3\text{OF}$  in both the gaseous and Ar matrix Raman spectra. We

assign this band to  $2 \delta_{\text{COF}}$  in Fermi resonance with  $\nu_{\text{OF}}$ . Using the method developed by Dixon (36) and used by Saier, Cousins and Basila (37), we calculate that a Fermi resonance perturbation of only  $1\text{--}3 \text{ cm}^{-1}$  would be necessary to give the intensity of the  $868 \text{ cm}^{-1}$  band relative to the  $882 \text{ cm}^{-1}$  band in Figure 3, if the  $868 \text{ cm}^{-1}$  band had zero intensity in the absence of Fermi resonance.

Both  $\text{CF}_3\text{OX}$  compounds have two bands in the IR and Raman spectra between  $200$  and  $300 \text{ cm}^{-1}$ . Initially it was assumed that both were  $\text{CF}_3$  rocking modes,  $\rho_{\text{CF}_3}$ , with the one in the  $A'$  block expected to be stronger in the Raman and the one in the  $A''$  block expected to be stronger in the IR.

The force constant calculations leave little doubt that the two rocking modes,  $\rho_{\text{CF}_3}$ , are far apart with the one in the  $A''$  block for  $\text{CF}_3\text{OX}$  under  $C_s$  symmetry being in the surprisingly high frequency range  $400\text{--}500 \text{ cm}^{-1}$ .

The microwave result (6) for  $\text{CF}_3\text{OF}$  requires the  $0 \rightarrow 1$  assignment in the  $\text{CF}_3$  torsion,  $\tau_{\text{CF}_3}$ , to the lowest frequency in the liquid Raman ( $144 \text{ cm}^{-1}$ ) and gas phase Raman ( $127 \text{ cm}^{-1}$ ) spectra for  $\text{CF}_3\text{OF}$  and by inference for  $\text{CF}_3\text{OCl}$  and other  $\text{CF}_3\text{OOX}$  series members also.

The choice of " $2 \tau_{\text{CF}_3}$ " and " $\rho_{\text{CF}_3}(A')$ " for the bands between  $200$  and  $300 \text{ cm}^{-1}$  is not unambiguous. If one band is above  $2 \times$  (" $\tau_{\text{CF}_3}$ "), and the other is slightly below  $2 \times$  (" $\tau_{\text{CF}_3}$ "), then the former is " $\rho_{\text{CF}_3}(A')$ " and the latter is



" $2 \tau_{\text{CF}_3}$ ". Such a comparison should be made with all frequencies from the same phase since the vapor torsion frequency is  $17 \text{ cm}^{-1}$  lower than the liquid frequency in the Raman spectra for  $\text{CF}_3\text{OF}$ . In general low frequency modes tend to increase in frequency upon condensation in contrast to stretching modes which tend to decrease in frequency upon condensation (38). Considering each phase separately, Table 9 was established to analyze the assignments of these two bands.

The frequency tabulation shown in Table 9 suggests 278 and  $233 \text{ cm}^{-1}$  as the correct choice for " $\rho_{\text{CF}_3}(\text{A}')$ " for gaseous  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$ , respectively. However, comparison across the " $\rho_{\text{CF}_3}(\text{A}')$ " row has the gaseous frequencies at higher values than the condensed phase frequencies contrary to normal expectations. It is, of course, possible that the nature of the normal mode could account for the frequencies being higher in the gas. Another disturbing feature is that the  $252 \text{ cm}^{-1}$  band is more intense than the  $278 \text{ cm}^{-1}$  band in the gaseous IR spectrum of  $\text{CF}_3\text{OF}$ . It is possible that the  $278 \text{ cm}^{-1}$  band is a part of the contour of a " $\rho_{\text{CF}_3}(\text{A}')$ " mode between 220 and  $300 \text{ cm}^{-1}$  with a center near  $265 \text{ cm}^{-1}$  and superposition of " $2 \tau_{\text{CF}_3}$ " near  $250 \text{ cm}^{-1}$  causes an apparent maximum at  $252 \text{ cm}^{-1}$ . The Raman gas bands (5) of  $\text{CF}_3\text{OF}$  at 272 and  $246 \text{ cm}^{-1}$  are of equal intensity and could be the contour of a single band. In the Raman spectra in an Ar matrix, only one band occurs for both  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$ . We assume that

Table 9. Frequency Comparison for Bands in the 200-300  $\text{cm}^{-1}$  Region for  $\text{CF}_3\text{OX}$  Compounds

	$\text{CF}_3\text{OF}$				$\text{CF}_3\text{OCl}$			
	Raman gas	Raman liquid	Raman Ar matrix	IR gas	Raman gas	Raman liquid	Raman Ar matrix	IR gas
$\nu_{\text{CF}_3}(\text{A}')$	272	259	256	278	---	220	239	233
$2\nu_{\text{CF}_3}(\text{A}')$	246	285?	---	252	---	233?	---	220
$\nu_{\text{CF}_3}(\text{A}'')$	127	144	---	(127) <sup>a</sup>	---	122	---	(108) <sup>b</sup>

a. Since we can't observe IR bands below about  $160\text{ cm}^{-1}$  on our Perkin-Elmer Model 180 spectrophotometer, we must use gas phase Raman frequencies or calculated values. Using the gas phase Raman data of Ref. 5 and our liquid Raman data for  $\text{CF}_3\text{OF}$  as reference, we get an 11.8% frequency shift from Raman liquid frequency to Raman gas frequency for the torsional mode.

b. Applying the same 11.8% frequency shift to  $\text{CF}_3\text{OCl}$  Raman liquid data at  $122\text{ cm}^{-1}$  we can get the vapor frequency of the torsional mode of  $\text{CF}_3\text{OCl}$  at  $108\text{ cm}^{-1}$ . This computational process is shown below:

Compound	Raman liquid	Raman gas	Shift	% shift
$\text{CF}_3\text{OF}$	$144\text{ cm}^{-1}$	$127\text{ cm}^{-1}$	$17\text{ cm}^{-1}$	11.8
$\text{CF}_3\text{OCl}$	$122\text{ cm}^{-1}$	$108\text{ cm}^{-1}$	$14\text{ cm}^{-1}$	11.8



band is " $\nu_{\text{CF}_3}(\text{A}')$ ". However, since the Ar matrix is too highly scattering to permit close enough approach to the Rayleigh line to observe " $\nu_{\text{CF}_3}$ " there is no opportunity to confirm that the single Raman band is above " $2 \nu_{\text{CF}_3}$ " and is most probably " $\nu_{\text{CF}_3}(\text{A}')$ " rather than " $2 \nu_{\text{CF}_3}$ ". Since matrix shifts from the gas are often only a few  $\text{cm}^{-1}$ , the single matrix frequency assumed to be " $\nu_{\text{CF}_3}(\text{A}')$ " matches best with 252 (or 265)  $\text{cm}^{-1}$  for the  $\text{CF}_3\text{OF}$  gaseous IR spectrum.

For gaseous  $\text{CF}_3\text{OCl}$  one can justify the use of 233  $\text{cm}^{-1}$  as " $\nu_{\text{CF}_3}(\text{A}')$ " on the grounds that it is more intense than the 220  $\text{cm}^{-1}$  absorption which is a shoulder in the IR of the gas. Also 233  $\text{cm}^{-1}$  is closer to the Ar matrix frequency which usually approximates gas frequencies more closely than liquid frequencies, we assume a single band in the Raman spectrum in the Ar matrix is " $\nu_{\text{CF}_3}(\text{A}')$ ".

For  $\text{CF}_3\text{OF}$  the Q branch at 429  $\text{cm}^{-1}$  has a shoulder at 431  $\text{cm}^{-1}$  which we had not tried to interpret. We can now suggest this as the " $\nu_{\text{CF}_3}(\text{A}'')$ " frequency on the basis of the normal coordinate analysis. The Raman spectra of Smardzewski and Fox (5,19) for an Ar matrix shows a single band at 433  $\text{cm}^{-1}$  whose width is comparable to that of other bands in the Ar matrix. In our opinion their Raman spectra can neither conform nor deny the possible presence of two fundamentals separated by a few  $\text{cm}^{-1}$  here. The liquid Raman band at 436  $\text{cm}^{-1}$  taken to be the counterpart of the band at

429  $\text{cm}^{-1}$  in the IR of the gas is polarized and should be " $\delta_{\text{COF}}$ ", an  $A'$  mode. This 436  $\text{cm}^{-1}$  band is slightly asymmetric to low frequencies so a weak depolarized band a few  $\text{cm}^{-1}$  away from 436  $\text{cm}^{-1}$  could possibly be present. If one accepts the " $\tau_{\text{CF}_3}$ ", " $2\tau_{\text{CF}_3}$ ", " $\rho_{\text{CF}_3}(A')$ " interpretation of the three bands between 100 and 300  $\text{cm}^{-1}$  and the general validity of our force field, then 431  $\text{cm}^{-1}$  is the only candidate for the " $\rho_{\text{CF}_3}(A'')$ " mode.

For  $\text{CF}_3\text{OCl}$ , there is no evidence of a band at 348  $\text{cm}^{-1}$  in either the Raman or IR spectra. The band with a Q branch at 393  $\text{cm}^{-1}$  has a small peak at 389  $\text{cm}^{-1}$  on its P branch which we assign to  $\nu_4$  of the impurity  $\text{SiF}_4$  whose presence is confirmed by its  $\nu_3$  at 1032  $\text{cm}^{-1}$  (39). We don't believe that both 393 and 389  $\text{cm}^{-1}$  are two nearly degenerate modes of  $\text{CF}_3\text{OCl}$  as we assigned for 429 and 431  $\text{cm}^{-1}$  in  $\text{CF}_3\text{OF}$ . The liquid Raman band at 397  $\text{cm}^{-1}$ , taken as the counterpart of the band at 393  $\text{cm}^{-1}$  in the IR of the gas, is slightly asymmetric to low frequencies. This could be the natural shape of the band or there could be a small contribution from  $\nu_4$  of  $\text{SiF}_4$  (a weak Raman band)(39). The shapes of the Raman bands at 436  $\text{cm}^{-1}$  in liquid  $\text{CF}_3\text{OF}$  and 397  $\text{cm}^{-1}$  in liquid  $\text{CF}_3\text{OCl}$  are sufficiently different that, in our opinion, it is possible that  $\text{CF}_3\text{OF}$  could have 2 bands present and  $\text{CF}_3\text{OCl}$  could have only one band present or one  $\text{CF}_3\text{OCl}$  band and one weak  $\text{SiF}_4$  band present. Since the 348  $\text{cm}^{-1}$  frequency in our calculation is in the  $A'$  block it is most reasonably

associated with the  $393\text{ cm}^{-1}$  IR band which must be in the A' block as it's Raman counterpart at  $397\text{ cm}^{-1}$  is polarized. Then the missing band for " $\rho_{\text{CF}_3}(\text{A}'')$ " is near  $420\text{ cm}^{-1}$  for our calculation in Table 7 and between  $420$  and  $465\text{ cm}^{-1}$  for some other calculations we have made. Our Raman data for  $\text{CF}_3\text{OCl}$  show a very weak band at  $430\text{ cm}^{-1}$  in every spectrum where the sensitivity is set high enough to record the  $611\text{ cm}^{-1}$  band, which is the weakest Raman line assigned to a fundamental " $\delta_{\text{CF}_3}(\text{A}'')$ ". The baseline in the IR is not flat in this region due to problems with exact compensation of a polyethylene band in the gas cell windows with two polyethylene sheets in the reference beam. It is difficult to be sure whether there is a very weak band near  $430\text{ cm}^{-1}$  in the IR or not. The fact that the band, if present, is weak compared to " $\rho_{\text{CF}_3}(\text{A}')$ " at  $233\text{ cm}^{-1}$  seems a little surprising for " $\rho_{\text{CF}_3}(\text{A}'')$ "; however, it is the only candidate for missing " $\rho_{\text{CF}_3}(\text{A}'')$ " mode.

Initially our sample of  $\text{CF}_3\text{Cl}$  contained a large amount of chlorine and the band in  $\text{CF}_3\text{OCl}$  at  $561\text{ cm}^{-1}$  in the Raman spectrum of the liquid was only visible as a small shoulder on the intense band due to the chlorine. Purification of the  $\text{CF}_3\text{OCl}$  sample gave the Raman spectra shown in Figures 7 and 8. The band at  $548\text{ cm}^{-1}$  in Figure 7 and 8 does not appear in the IR spectrum of gaseous  $\text{CF}_3\text{OCl}$ . Smardzewski and Fox (5,19) observe bands in Raman spectra

of  $\text{CF}_3\text{OCl}$  in an Ar matrix at  $547$  and  $539\text{ cm}^{-1}$  with relative intensities of 9 and 10, respectively. Our band at  $548\text{ cm}^{-1}$  in the Raman spectrum of liquid  $\text{CF}_3\text{OCl}$  appears to consist of two overlapping bands of equal intensity at  $\sim 551$  and  $\sim 545\text{ cm}^{-1}$ . Since all the  $\text{CF}_3\text{OCl}$  fundamentals are reasonably accounted for, the band(s) at  $\sim 548\text{ cm}^{-1}$  must be either a non fundamental of  $\text{CF}_3\text{OCl}$  or an impurity.

The assignment of the  $548\text{ cm}^{-1}$  band in the liquid Raman spectrum of  $\text{CF}_3\text{OCl}$  to a difference band is reasonable on a frequency basis but not on the basis of intensity and polarization. Using liquid Raman frequencies " $\delta_{\text{CF}_3}(\text{A}')$ " - " $\tau_{\text{CF}_3}$ " =  $666 - 122 = 544\text{ cm}^{-1}$  in good agreement with  $548\text{ cm}^{-1}$ . However, the transition from ( $v_{12}=1$ , all other  $v_i=0$ , an A" state) to ( $v_5=1$ , all other  $v_i=0$ , an A' state) is a non totally symmetric transition and should be depolarized. The  $548\text{ cm}^{-1}$  band is polarized. The population of ( $v_{12}=1$ , all other  $v_i=0$ ) is 10-17 % of that of (all  $v_i=0$ ) for  $v_{12}=\tau_{\text{CF}_3}=122\text{ cm}^{-1}$  between  $77$  and  $100^\circ\text{K}$  (estimated range of sample temperature with liquid  $\text{N}_2$  in the cold bath of the low temperature cell). The fact that the  $548\text{ cm}^{-1}$  band has 80% of the intensity of the " $\delta_{\text{CF}_3}(\text{A}')$ " fundamental at  $561\text{ cm}^{-1}$  is difficult to explain for a non fundamental with no apparent Fermi resonance possibilities and such an unfavorable Boltzmann population ratio. On both the intensity and polarization basis we feel that the  $548\text{ cm}^{-1}$  band must be due to a species other than

$\text{CF}_3\text{OCl}$ .

Possible impurities in the  $\text{ClF}$  used to prepare  $\text{CF}_3\text{OCl}$  by addition to  $\text{COF}_2$  are  $\text{ClF}_5$  and  $\text{ClF}_3$ . However, the Raman spectra of liquid  $\text{ClF}_5$  (40) and liquid  $\text{ClF}_3$  (41) are such that both compounds would be detected by intense bands well separated from any of the  $\text{CF}_3\text{OCl}$  bands in Figures 7 and 8. The source of the  $548\text{ cm}^{-1}$  band(s) must be so similar to  $\text{CF}_3\text{OCl}$  that the two cannot be easily separated by physical means. Another possibility is an additional reaction product from the preparation.

For the case of  $\text{CF}_3\text{OF}$  prepared by addition of  $\text{F}_2$  to  $\text{COF}_2$ ,  $\text{CO}_2$  impurity in the  $\text{COF}_2$  leads to the production of some  $\text{CF}_2(\text{OF})_2$ . By analogy the addition of  $\text{ClF}$  to  $\text{CO}_2$  might lead to a variety of materials of the general formula  $\text{CF}_w\text{Cl}_x(\text{OF})_y(\text{OCl})_z$ . Here  $w, x, y, z$  may have the values 0, 1, or 2 in various combinations subject to the restrictions that  $(w+x) = 2$  and  $(y+z) = 2$ . For compounds with C-Cl bonds ( $x \neq 0$ ) and compounds with O-F bonds ( $y \neq 0$ ), Raman bands should be present that are well separated from any of the  $\text{CF}_3\text{OCl}$  bands in Figures 7 and 8. One that might be less easily detected is  $\text{CF}_2(\text{OCl})_2$ . It is possible to estimate the frequencies for  $\text{CF}_2(\text{OCl})_2$  from those of  $\text{CF}_3\text{OCl}$  by using differences between  $\text{CF}_3\text{OF}$  and  $\text{CF}_2(\text{OF})_2$  frequencies. These estimates suggest the  $\text{CF}_2(\text{OCl})_2$  might have a strong band at  $548\text{ cm}^{-1}$  but not no traces of other bands in addition

to those in Figures 7 and 8.

An additional possibility is further reaction of  $\text{CF}_3\text{OCl}$  to form  $\text{CF}_3\text{OClF}_2$ . The  $\text{CF}_3\text{O}$  fragments of both  $\text{CF}_3\text{OCl}$  and  $\text{CF}_3\text{OClF}_2$  might have indistinguishable spectra and differentiation would depend on vibrations of the  $\text{ClF}_2$  fragment. Since  $\text{ClF}_3$  has a  $\text{ClF}$  stretching mode at  $529\text{ cm}^{-1}$  (gas phase, with liquid at lower frequency), the  $\text{ClF}_2$  fragment of  $\text{CF}_3\text{OClF}_2$  could be responsible for the  $548\text{ cm}^{-1}$  band. However, it is difficult to believe that the remaining modes of the  $\text{ClF}_2$  fragment would not produce additional bands in Figures 7 and 8.

Chlorine dissolved in  $\text{CF}_3\text{OCl}$  would be expected to give a band (really three bands for  $^{35}\text{Cl}_2$ ,  $^{35}\text{Cl}^{37}\text{Cl}$  and  $^{37}\text{Cl}_2$ ) near  $545\text{ cm}^{-1}$ . Liquid chlorine in our low temperature cell with liquid nitrogen in the bath gave bands at 547, 540, and  $533\text{ cm}^{-1}$  in the intensity ratio 6.5 : 4.5 : 1 (theoretical 9 : 6 : 1). Condensation of chlorine into a  $\text{CF}_3\text{OCl}$  sample of purity similar to Figures 7 and 8 gave bands at 552 and  $544\text{ cm}^{-1}$  and a possible shoulder at  $537\text{ cm}^{-1}$ . The  $548\text{ cm}^{-1}$  band in  $\text{CF}_3\text{OCl}$  cannot be due to chlorine in the same environment as pure liquid chlorine or excess chlorine in  $\text{CF}_3\text{OCl}$ . However, it would be possible for  $\text{CF}_3\text{OCl}$  to form a complex with chlorine of sufficient strength that it is not possible to remove all the chlorine from  $\text{CF}_3\text{OCl}$ . Then the  $548\text{ cm}^{-1}$  band could be due to the stretching of the  $\text{Cl}-\text{Cl}$  bond in the



complex.

The observation of this band in the Raman spectra of liquid  $\text{CF}_3\text{OCl}$  near liquid nitrogen temperature and of  $\text{CF}_3\text{OCl}$  in an Ar matrix at  $8^\circ\text{K}$  but not in the IR spectrum of the gas at room temperature is reasonable. Low temperature favors complex formation and the complex is probably not present in the gas phase at room temperature. Even if the complex were present in the gas it would probably be too weak to cause the IR inactive stretching of gaseous  $\text{Cl}_2$  to become observable in the IR spectrum of the gaseous complex.

The proposed complex of chlorine with  $\text{CF}_3\text{OCl}$  might be expected to show isotope splitting in its chlorine stretching mode. Assuming the complexation takes place via the chlorine atom of  $\text{CF}_3\text{OCl}$  (analogous to the  $\text{X}_3^-$  halide complex ions like  $\text{I}_3^-$ ), eight complexes are possible as follows:

No.	Mass Arrangement $\text{CF}_3\text{OCl} \dots \text{Cl} \text{ --- } \text{Cl}$			Relative Abundance
1	35	35	35	27
2	35	35	37	9
3	35	37	35	9
4	35	37	37	3
5	37	35	35	9
6	37	35	37	3
7	37	37	35	3
8	37	37	37	1

The effect of the mass of the chlorine atom in  $\text{CF}_3\text{OCl}$  on the frequency for stretching the Cl-Cl bond from  $\text{Cl}_2$  in the complex depends on the strength of the complex. The limit for a weak complex could be three bands due to three degenerate sets as follows: 1 and 5 of relative intensity 36; 2, 3, 6, and 7 of relative intensity 24; and 4 and 8 of relative intensity 4. As complex strength increases these degeneracies would be broken as the mass of the chlorine atom in  $\text{CF}_3\text{OCl}$  begins to influence the frequency. A reasonable extreme would be where the vibration is still best treated as a perturbed diatomic molecule stretching rather than a three body problem with antisymmetric and symmetric skeletal stretching and skeletal bending. However, the degeneracies are broken and the eight frequencies might tend to group into the following four bands: 1 of relative intensity 27; 2, 3, and 5 of relative intensity 27; 4, 6, and 7 of relative intensity 9; and 8 of relative intensity 1.

For the weak extreme, the third band probably would not be observed as it is too weak so a higher frequency band 1.5 times as intense as a lower frequency band is to be expected. For the strong extreme the fourth band would definitely be too weak to observe and the extent of overlap of the other three bands is uncertain. A likely result would be that the third band of intensity 9 would overlap the second band of intensity 27 to give a single asymmetric band or a



band with a low frequency shoulder. This superposition would result in a lower frequency band with relative intensity 27 (or more from overlap of the third band) and a higher frequency band of relative intensity 27 (from complex 1). Thus, two resolvable bands seem reasonable for the range of complex strengths suggested. The higher frequency band would be expected to be from 1.5 times as intense to slightly less intense than the lower frequency band depending on the strength of the complex.

The formation of a  $\text{CF}_3\text{OCl} \dots \text{Cl}-\text{Cl}$  complex will then account for a band at  $548 \text{ cm}^{-1}$  in the Raman spectrum of liquid  $\text{CF}_3\text{OCl}$  appearing to consist of two overlapping bands of equal intensity at  $551$  and  $545 \text{ cm}^{-1}$ . The observation by Smardzewski and Fox (5,19) of bands at  $547$  and  $539 \text{ cm}^{-1}$  with relative intensities of 9 and 10, respectively, in the Raman spectrum of  $\text{CF}_3\text{OCl}$  in an Ar matrix is also consistent. The conditions of a liquid near  $77^\circ\text{K}$  and an Ar matrix at  $8^\circ\text{K}$  are conditions that should favor complex formation. The improved resolution in the Ar matrix is also reasonable.

This discussion is, of course, speculative and does not prove anything. However, this explanation does account for the fact that no additional unexplained bands are found. We believe it is unreasonable to associate the  $548 \text{ cm}^{-1}$  band with pure  $\text{CF}_3\text{OCl}$ . Thus some other species must be responsible. At present, the  $\text{CF}_3\text{OCl} \dots \text{Cl}-\text{Cl}$  complex provides the

simplest explanation and we consider it the most likely cause of the  $\sim 548 \text{ cm}^{-1}$  band.

### C. Comments on the $L^{-1}$ Matrices and the Normal Coordinates

The matrix  $L^{-1}$  (eigenvector inverse matrix) gives the transformation between symmetry and normal coordinates:  $Q = L^{-1}S$ . These transformation matrices for both the  $A'$  and  $A''$  symmetry species of  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  are shown in Tables 10 and 11, respectively, for the calculation with  $(1/\lambda_i)^2$  weighting. Thus,  $Q_1$  essentially equals approximately  $2 S_1$  and  $Q_9$  approximately  $2 S_{10}$  for both molecules. Thus they are described as the antisymmetric stretches of the  $\text{CF}_3$  group for the  $A'$  and  $A''$  symmetry species, respectively. In general the  $L^{-1}$  matrices in Tables 10 and 11 show a large amount of mixing of symmetry coordinates to be present in the normal coordinates. For example in  $\text{CF}_3\text{OF}$  we have  $Q_3(947 \text{ cm}^{-1}) \cong 3.39 S_2 + 1.63 S_7$  and in  $\text{CF}_3\text{OCl}$   $Q_3(919 \text{ cm}^{-1}) \cong 3.73 S_2 + 1.82 S_7$ . Thus, the normal coordinate  $Q_3$ , assigned as the CO stretching mode, has important contributions from both symmetrical  $\text{CF}_3$  stretching and the CO stretching motions. This situation is not unreasonable when one considers the relative masses of the carbon and three fluorine atoms of the  $\text{CF}_3$  group. A stretching of the C-O bond would tend to shorten the C-F bonds in a symmetrical manner. The resemblance of the form of normal coordinates called CO stretching mode

Table 10. Symmetrized  $L^{-1}$  Matrix for  $CF_3OF^a$ 

	$S_1$	$S_2$	$S_3$	$S_4$	$S_5$	$S_6$	$S_7$	$S_8$	$S_9$	$S_{10}$	$S_{11}$	$S_{12}$	$S_{13}$
$Q_1$	<u>1.69</u>	-0.18	0.26	0.28	0.16	-0.44	0.66	-0.14					
$Q_2$	0.90	<u>1.07</u>	-0.51	0.12	-0.55	-0.02	<u>-1.26</u>	-0.03					
$Q_3$	-0.80	<u>3.39</u>	0.84	0.00	0.08	-0.47	<u>1.63</u>	-0.18					
$Q_4$	-0.71	<u>-1.44</u>	<u>2.18</u>	0.15	-0.36	-0.30	-0.81	-0.46					
$Q_5$	-1.43	<u>-1.67</u>	<u>-1.77</u>	0.57	<u>-1.77</u>	-1.15	1.30	-0.32					
$Q_6$	<u>-2.45</u>	0.45	-0.76	<u>2.45</u>	1.09	-0.06	-1.29	-0.62					
$Q_7$	0.74	0.09	<u>2.25</u>	1.67	-1.45	<u>2.04</u>	<u>2.44</u>	<u>1.63</u>					
$Q_8$	0.84	-0.29	-0.07	0.01	1.18	<u>-3.97</u>	-2.85	<u>3.73</u>					
$Q_9$										<u>2.11</u>	0.32	-0.15	0.00
$Q_{10}$										<u>-2.91</u>	<u>2.35</u>	-1.19	0.02
$Q_{11}$										0.55	1.84	<u>2.58</u>	-0.07
$Q_{12}$										0.87	0.27	<u>2.13</u>	<u>1.29</u>

a. See footnote a in Table 6 for numbering system for  $S_i$  and  $Q_i$ .

Table 11. Symmetrized  $L^{-1}$  Matrix for  $CF_3OCl^a$ 

	$S_1$	$S_2$	$S_3$	$S_4$	$S_5$	$S_6$	$S_7$	$S_8$	$S_9$	$S_{10}$	$S_{11}$	$S_{12}$	$S_{13}$
$Q_1$	<u>1.87</u>	-0.06	0.06	0.29	0.10	-0.38	0.40	-0.14					
$Q_2$	0.46	<u>1.01</u>	0.01	0.07	-0.61	-0.05	<u>-1.37</u>	-0.05					
$Q_3$	-0.27	<u>3.73</u>	-0.29	-0.05	0.21	-0.17	<u>1.82</u>	0.03					
$Q_4$	<u>-1.68</u>	0.20	<u>1.94</u>	0.43	0.09	-0.96	0.09	-0.77					
$Q_5$	-1.00	-1.63	-0.93	0.45	<u>-2.10</u>	-0.74	<u>1.83</u>	-0.21					
$Q_6$	<u>-2.19</u>	0.23	-1.09	<u>2.67</u>	0.70	0.26	-0.96	-0.26					
$Q_7$	-1.26	0.23	<u>3.52</u>	1.30	-1.29	1.63	<u>2.39</u>	<u>2.29</u>					
$Q_8$	0.55	-0.43	-1.37	-0.38	1.42	<u>-5.61</u>	<u>-3.14</u>	<u>3.88</u>					
$Q_9$									<u>2.10</u>	0.32	-0.15	0.00	
$Q_{10}$									<u>-2.90</u>	<u>2.40</u>	-1.12	0.02	
$Q_{11}$									0.63	1.79	<u>2.59</u>	-0.08	
$Q_{12}$									0.88	0.18	<u>2.58</u>	<u>1.50</u>	

a. See footnote a in Table 6 for numbering system for  $S_i$  and  $Q_i$ .

in both  $\text{CF}_3\text{OX}$  molecules tends to support the correctness of the new assignment of this mode. It can be seen that the chief contribution to the normal coordinates came from the related symmetry coordinates in some cases. Then the word description of the mode such as 'antisymmetric  $\text{CF}_3$  stretching' is still qualitatively correct. However some modes involve extensive mixing of the symmetry coordinates and the word descriptions do not have even qualitative significance.

#### D. Comments on the Potential Energy Distribution

Another method of describing molecular vibrations is through the potential energy distribution (PED) which is obtained from the L matrix (eigenvector matrix). The PED provides a convenient means for representing the contributions to the potential energy of the system from the symmetry or internal coordinate.

Tables 12 and 13 show the degree of localization of a particular normal mode coordinate among the various symmetry force constants. In general, the results parallel those of the  $L^{-1}$  matrices in the behavior of mixing, but the distributions show much greater localization than do the  $L^{-1}$  matrices. The most conspicuous examples are from the torsional modes of both molecules. In this case about 95% of the contribution is from the torsional force constant.

Table 12. Potential Energy Distribution<sup>a</sup> for CF<sub>3</sub>OF

$\nu_i$	F <sub>1</sub>	F <sub>2</sub>	F <sub>3</sub>	F <sub>4</sub>	F <sub>5</sub>	F <sub>6</sub>	F <sub>7</sub>	F <sub>8</sub>	F <sub>10</sub>	F <sub>11</sub>	F <sub>12</sub>	F <sub>13</sub>
1	1321	<u>76</u>		16	5	11	12	10				
2	1214	17	38		38		<u>54</u>					
3	941	5	<u>60</u>	7		8	8					
4	883		<u>76</u>		5	5	18	7				
5	674	6	12		<u>36</u>	17	11					
6	589	5		<u>61</u>	12							
7	420		8	24	10	21	5	<u>29</u>				
8	267					38	6	<u>58</u>				
9	1256								<u>105</u>	20	12	
10	616								8	<u>55</u>	15	
11	429									31	<u>76</u>	
12	128											<u>96</u>

a. Contributions of less than 5% are not included in this table. Numbering of symmetrized force constants F<sub>i</sub> follows the numbering of symmetry coordinates in Table 4. No F<sub>9</sub> appears because S<sub>9</sub> is the redundant symmetry coordinate in the A' symmetry block. The numbering of frequencies is from Table 6. See footnote a in Table 6 for the numbering system for  $\nu_i$  and S<sub>i</sub>.

Table 13. Potential Energy Distribution<sup>a</sup> for CF<sub>3</sub>OC1

$\nu_i$	$F_1$	$F_2$	$F_3$	$F_4$	$F_5$	$F_6$	$F_7$	$F_8$	$F_{10}$	$F_{11}$	$F_{12}$	$F_{13}$
1 1309	<u>89</u>			18		9	5	10				
2 1208	5	41			46		<u>67</u>					
3 935		<u>65</u>					16					
4 774	9		<u>51</u>			22		14				
5 658		5	9		<u>48</u>	8	15					
6 571			8	<u>73</u>	5							
7 348			30	12	6	9		<u>38</u>				
8 217						<u>50</u>		44				
9 1262									<u>104</u>	20	11	
10 613									8	<u>57</u>	13	
11 422										30	<u>77</u>	
12 108											5	<u>96</u>

a. Contribution of less than 5% are not included in this table. Numbering of symmetrized force constants  $F_i$  follows the numbering of symmetry coordinates in Table 4. No  $F_9$  appears because  $S_9$  is the redundant symmetry coordinate in the A' symmetry block. The numbering of frequencies is from Table 7. See footnote a in Table 6 for the numbering system for  $\nu_i$  and  $S_i$ .

## E. Comments on the Barrier to Internal Rotation

It is possible to calculate the height of the threefold barrier hindering internal rotation in  $\text{CF}_3\text{OX}$  molecules from the proposed geometry listed in Table 3 and from the observed torsional frequencies. Since the torsional frequencies were below the low frequency limit of the Perkin-Elmer Model 180 infrared spectrophotometer in our laboratory, their values were obtained from the gas phase and liquid Raman data which are discussed in Table 9.

The frequencies  $144\text{ cm}^{-1}$ ,  $122\text{ cm}^{-1}$ ,  $127\text{ cm}^{-1}$ , and  $108\text{ cm}^{-1}$  seem unexpectedly low for rocking vibrations and unexpectedly high for torsions. Eleven of the twelve fundamentals for a  $\text{CF}_3\text{OX}$  structure are reasonably assigned to Raman bands at frequency shifts greater than  $200\text{ cm}^{-1}$  for both  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$ . These four frequencies must then be assigned to the  $\text{CF}_3$  torsion mode, but the question is open as to whether the transitions responsible for the bands have  $\Delta v = 1$  or  $\Delta v = 2$ . For a  $\text{CF}_3\text{OX}$  structure of  $C_s$  symmetry, as deduced from electron diffraction for  $\text{CF}_3\text{OF}$  (13),  $\Delta v = 2$  transitions are totally symmetric while  $\Delta v = 1$  transitions are non-totally symmetric for a non-totally symmetric mode such as the  $\text{CF}_3$  torsion. Observation of a small depolarization ratio would eliminate a  $\Delta v = 1$  transition while a band which is too weakly polarized to be distinguished from being depolarized could be either a  $\Delta v = 1$  or  $\Delta v = 2$  transition.



For liquid  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  where depolarization ratios were measured for the  $144\text{ cm}^{-1}$  and  $122\text{ cm}^{-1}$  bands, a definite distinction between the bands being weakly polarized and depolarized could not be made.

The microwave spectrum of  $\text{CF}_3\text{OF}$  has been analyzed by Buckley and Weber (6). Vibrational satellites were observed at  $194^\circ\text{K}$  and the intensity ratio of the satellite line to the ground state line was about 0.4. This ratio leads to a torsional barrier of 3.9 kcal/mole in reasonable agreement with 4.4 kcal/mole using the  $0 \rightarrow 1$  assignment for the  $127\text{ cm}^{-1}$  band in gaseous  $\text{CF}_3\text{OF}$ . Working in the opposite direction, the gaseous frequency of  $127\text{ cm}^{-1}$  for  $\text{CF}_3\text{OF}$  at  $194^\circ\text{K}$  gives a vibrational state population ratio of 0.39 in excellent agreement with the observed ratio of 0.4. If the  $127\text{ cm}^{-1}$  band were  $0 \rightarrow 2$  in the  $\text{CF}_3$  torsion then  $0 \rightarrow 1$  would be at  $65\text{ cm}^{-1}$  giving a vibrational state population ratio of 0.62 at  $194^\circ\text{K}$ . The difference in the ratios of 1.6 (i.e.,  $0.62/0.39$ ) seems to exceed any possible experimental error in measuring the intensity ratio. Once the decision is made that the observed Raman band is the  $0 \rightarrow 1$  transition, then this frequency is a more reliable source of the barrier height on the grounds that a frequency should be more accurately known than an intensity ratio. Thus we would propose the best value of the barrier for the  $\text{CF}_3$  torsion in  $\text{CF}_3\text{OF}$  as 4.4 kcal/mole. For  $\text{CF}_3\text{OCl}$  assuming the same

percentage frequency shift on phase change as  $\text{CF}_3\text{OF}$ , the liquid Raman frequency of  $122\text{ cm}^{-1}$  becomes  $108\text{ cm}^{-1}$  for the vapor and the barrier for  $\text{CF}_3$  torsion is  $4.2\text{ kcal/mole}$ .

The barrier heights,  $V_3$ , of the  $\text{CF}_3$  rotor were calculated following the procedures developed by Fateley and Miller (42-44). The parameters used for this calculation are listed in Table 14. Agreement between the two values of the barrier height in the two molecules in the same phase is good. These values are slightly higher than the microwave result of  $3.9\text{ kcal/mole}$  (13). A band at  $56\text{ cm}^{-1}$  was assigned as the  $0 \rightarrow 1$  transition of the  $\text{CF}_3$  torsion and a potential barrier of  $395\text{ cm}^{-1}$  ( $1.13\text{ kcal/mole}$ ) was determined with an assumed geometry by Wilt and Jones (4). The existence of this band is questionable based on our IR and Raman data and the Raman data of Smardzewski and Fox (5). This strikingly low value for the barrier height is also inconsistent with the microwave and electron diffraction results.

Table 14. Summary of Parameters in Barrier Height Calculation

Parameter	CF <sub>3</sub> OF		CF <sub>3</sub> OCl	
	Raman gas	Raman liquid	Raman gas	Raman liquid
$\nu_{CF_3}$ (cm <sup>-1</sup> )	127	144	108 <sup>a</sup>	122
$I_\alpha$ (amu-Å <sup>2</sup> ) <sup>b</sup>	88.06	88.06	88.06	88.06
$rI_\alpha$ (amu-Å <sup>2</sup> ) <sup>b</sup>	13.85	13.85	18.40	18.40
$\Delta b_{\nu\sigma}$ <sup>c</sup>	46.37	52.58	52.40	59.19
$S^c$	560.8	718.3	713.4	906.3
$F$ (amu-Å <sup>2</sup> )	1.217	1.217	0.916	0.916
$V_3$ (kcal/mole) <sup>c</sup>	4.4	5.6	4.2	5.3

- a. Calculated assuming the same percentage frequency shift on phase change as CF<sub>3</sub>OF.
- b. The quantity  $rI_\alpha$ , the reduced moment for internal rotation, is given by

$$rI_\alpha = (1 - I_\alpha \sum_{i=1}^3 \lambda_i / I_i) I_\alpha$$

here  $I$  is the moment of inertia of the CF<sub>3</sub> group about its symmetry axis,  $I_i$  and  $\lambda_i$  are the principal moments of inertia of the entire molecule and direction cosines of the CF<sub>3</sub> symmetry axis with respect to the principal axes, respectively.

- c. The equations used to calculate  $V_3$  (cm<sup>-1</sup>) are as follows:

$$\nu_{CF_3} = \Delta E_{\nu\sigma} = (2.25) F \Delta b_{\nu\sigma}$$

$$F = 16.8576 / rI_\alpha \text{ (amu-Å}^2\text{)}$$

$$V_3 = 2.25 FS \text{ (cm}^{-1}\text{)}$$

The Mathieu function table of  $b_{\nu\sigma}$  (the eigenvalue of the Mathieu equation) VS.  $S$  (the dimensionless parameter) used is from reference (44).

II. RAMAN SPECTRA OF  $\text{CF}_2(\text{OF})_2$  AS A FUNCTION OF TEMPERATURECHAPTER I  
INTRODUCTION

Gady and co-workers were the first to report the preparation of a carbon-containing hypofluorite,  $\text{CF}_3\text{OF}$  (11). Since then a number of new carbon containing -OF compounds have been reported (45). Bis(fluoroxo)perfluoroalkanes are a new class of compounds coming from this development. Bis(fluoroxo)difluoromethane,  $\text{CF}_2(\text{OF})_2$ , is the parent member of this class (46). It is conveniently prepared in high yield by static, cesium fluoride catalyzed fluorination of carbon dioxide (47).

The longer known hypofluorite,  $\text{CF}_3\text{OF}$ , undergoes a variety of interesting reactions with both inorganic and organic reagents. The spectroscopic investigations and intramolecular force field calculation of this compound are reported in Part I of this study.

Because of the apparent similarity of  $\text{CF}_2(\text{OF})_2$  and  $\text{CF}_3\text{OF}$ , it was of interest to study the physical properties of this compound by using vibrational spectroscopy.

Structural investigation of  $\text{CF}_2(\text{OF})_2$  by the analysis of its fluorine nuclear magnetic resonance spectrum suggested that the two triplets (1:2:1) of equal area at -155.7 and +81.7 ppm (trichlorofluoromethane was the internal reference) be assigned to the two equivalent OF and CF fluorines, res-

pectively (46,47). A somewhat detailed infrared spectrum of  $\text{CF}_2(\text{OF})_2$  was reported by Mitchell and Merrit (48). Measured frequencies of the observed infrared bands (in  $\text{cm}^{-1}$ ) of this compound were presented. A possible  $\text{C}_{2v}$  symmetry with 15 fundamental vibrations was suggested.

During the course of this study, the concept of rotational isomerism was proposed to explain the observation of more bands in the low temperature liquid Raman spectrum of this molecule than what would be allowed by the selection rules for a single spatial structure. The additional bands could not assigned to overtone or combination modes and could only be satisfactorily explained on the basis of this compound possessing more than one geometric structure.

A subsequent study of the temperature effect on Raman spectra confirmed this proposition, as the spectra of this molecule were temperature sensitive. Certain bands became decidedly weaker and some became much stronger when the temperature of this sample was increased from near liquid nitrogen temperature ( $-196^\circ\text{C}$ ) up to above its boiling point ( $-40^\circ\text{C}$ ).

This observation may be rationalized in terms of a depopulation of a thermodynamically less stable isomeric structure with decreasing temperature. The molecules may exist as an equilibrium mixture of two or more rotational isomers or conformers over a temperature range. In the present

study we have assumed a two isomer mixture although the possibility of more than two isomers being present cannot be rigorously excluded.

The experimental determination of absolute Raman intensity was known to be very difficult. Consequently a relative intensity comparison was used to observe the population change.

The present report concerns preliminary work on this subject. The vibrational spectra of this compound were recorded and tentative interpretation is presented. More experimental data are needed to fully explore the dynamic nature of this conformational system. To this end some suggestions for further study involving other spectroscopic techniques are presented.

CHAPTER II  
EXPERIMENTAL

The samples of bis(fluoroxy)perfluoromethane,  $\text{CF}_2(\text{OF})_2$ , used in this study were prepared by Dr. D. D. DesMarteau at Kansas State University. The preparation was that of Hohorst and Shreeve (47), and involved the static fluorination of carbon dioxide with excess fluorine in the presence of cesium fluoride. The crude sample was trapped at liquid nitrogen temperature and then purified by gas chromatography. The purity of this sample was monitored by recording its infrared spectrum. The infrared spectrum of this sample shown in Figure 13 is similar to that reported by Thompson (46) and Mitchell and Merrit (48). The spectrum shown in Figure 14 does not appear to agree as well with that of Thompson (46) but is qualitatively similar to that of Mitchell and Merrit (48).

The infrared spectra were recorded using procedures described in the experimental section in Chapter II, Part I of this thesis. Liquid Raman spectra were recorded in a low temperature cell similar in design to that of Brown et.al. (18) on a system for Laser Raman spectroscopy consisting of a Spex Model 14018 Double Monochromator with a RCA C31034 photomultiplier tube and Princeton Applied Research Model 1140AC Quantum Photometer. Excitation was with the 5145 Å line of a Spectra-Physic Model 164-03 argon ion laser. A complete description of the system for Laser Raman spectroscopy appears elsewhere (8).



EXPLANATION OF FIGURE 13

Gas-phase Infrared Spectrum of  $\text{CF}_2(\text{OF})_2$   
in the Region  $3000\text{--}300\text{ cm}^{-1}$ .

Spectrometer: Perkin-Elmer Model 180

Sampling Method: 5 cm Path Length Cell  
with AgCl Windows

Pressure: 240 mm Hg

Energy Mode: Constant  $I_0$

Resolution:  $2.0\text{ cm}^{-1}$  at  $4000\text{ cm}^{-1}$

Gain: 6

Slit Program: 5

Time Constant: 1

Scan Time: Fine 5, Coarse 10


Suppression: 5

Temperature:  $25^\circ\text{C}$



By means of appropriate coolants, the low temperature cell was used to record the liquid Raman spectrum with the cell bath at four different temperatures of  $-40^{\circ}\text{C}$ ,  $-78^{\circ}\text{C}$ ,  $-135^{\circ}\text{C}$ , and  $-196^{\circ}\text{C}$ . Liquid nitrogen was used as coolant to get the bath temperature down to  $-196^{\circ}\text{C}$ . The  $-135^{\circ}\text{C}$  temperature was obtained by using a dry ice-freon 12-liquid nitrogen bath. Constant liquid nitrogen addition followed by vigorous stirring (to break the ice layer) was required to keep temperature within  $135 \pm 2^{\circ}\text{C}$ . A dry ice-freon 12 mixture was used to get the bath temperature within  $1^{\circ}\text{C}$  of  $-78^{\circ}\text{C}$ . A freon 12-liquid nitrogen bath was used for the temperature of  $-40 \pm 1^{\circ}\text{C}$ . The normal boiling point of  $\text{CF}_2(\text{OF})_2$  is  $-64^{\circ}\text{C}$  which is below the experimental bath temperature of  $-40^{\circ}\text{C}$ . After long irradiation with the incident laser beam, bubbles, which were probably caused by local heating with subsequent vigorous vaporization, come up during the recording. These bubbles may have produced some fluctuation in the scattered light. With the loss of sample through constant vaporization, the effect on signal intensities and depolarization ratios is hard to estimate.

The spectrum of the gas was obtained with the sample contained in a cell similar in design to the standard Spex multipass gas cell. High laser power and high gain were required for recording the gas phase Raman spectra. Due to this experimental difficulty, the gas phase Raman spectrum (Figure 15) was not as good in quality as the liquid Raman spectrum.



EXPLANATION OF FIGURE 15

Raman Spectrum of Gas-phase  $\text{CF}_2(\text{OF})_2$   
in the Region  $1000\text{--}800\text{ cm}^{-1}$

Spectrometer: Spex Model 14018

Slit Setting:  $200\text{ m}\mu$

Gain: 300

Filter: 2.2%

Scan Speed:  $0.5\text{ cm}^{-1}/\text{sec}$


Chart Speed:  $3\text{ cm}/\text{min}$


Source:  $\text{Ar}^+$  Laser at  $5145\text{ \AA}$

Power:  $1.4\text{ W}$

Temperature:  $25^\circ\text{C}$

Pressure:  $\sim 600\text{ mm Hg}$





### CHAPTER III RESULTS AND DISCUSSION

#### A. General

No detailed structural studies like electron diffraction or microwave spectroscopy have been published for the molecule  $\text{CF}_2(\text{OF})_2$ . A F-19 nuclear magnetic resonance study has indicated that this molecule has two equivalent -OF groups and that the two fluorine atoms of the  $-\text{CF}_2-$  group are equivalent as well (46,47). This NMR study was performed at ambient temperature using a 10% solution of this compound in trichlorofluoromethane, which also served as the internal reference. It is not clear what the influence of the solvent on the conformer equilibrium might be.

An inspection of the Raman spectrum of liquid  $\text{CF}_2(\text{OF})_2$  in Figure 16 shows at least 22 well-defined Raman bands with some additional shoulders. If one assumes only one molecular configuration, either with  $\text{C}_{2v}$  symmetry as indicated by the NMR study or  $\text{C}_s$  symmetry as compared with  $\text{CF}_3\text{OF}$ , then the maximum number of bands that should be observed is 15, which is the total number of normal vibrations for the  $\text{CF}_2(\text{OF})_2$  molecule. Difference bands can be ruled out because of the low bath temperature used to record the Raman data. One is then forced to look for combination or overtone bands to explain the "extra" Raman bands.

For example the  $524\text{ cm}^{-1}$  band could be explained as arising from the combination,  $243 + 280 = 523\text{ cm}^{-1}$ , but the  $524\text{ cm}^{-1}$  Raman band is as intense as the  $280\text{ cm}^{-1}$  band.

EXPLANATION OF FIGURE 16

Raman Spectrum of Liquid  $\text{CF}_2(\text{OF})_2$

Spectrometer: Spex Model 14018

Slit Setting: 200  $\mu\text{m}$

Gain: 3 K

Filter: 1.3%

Scan Speed: 1  $\text{cm}^{-1}/\text{sec}$

Chart Speed: 3  $\text{cm}/\text{min}$

Source:  $\text{Ar}^+$  Laser at 5145  $\text{\AA}$

Power: 300 mW

Temperature:  $-196^\circ\text{C}$

Therefore, it is unlikely to be a combination band on the basis of its intensity relative to the fundamentals from which it would have to arise. One is then forced on the basis of the Raman spectrum alone to consider a second rotational isomer or conformer as being present at  $-196^{\circ}\text{C}$ . Then, the additional Raman bands could arise from this second species. The effects of temperature on the Raman spectrum of  $\text{CF}_2(\text{OF})_2$  in the liquid state were observed at four different bath temperatures:  $-40^{\circ}\text{C}$ ,  $-78^{\circ}\text{C}$ ,  $-135^{\circ}\text{C}$  and  $-196^{\circ}\text{C}$ . In the present study, the intensities of the Raman bands at  $944$  and  $867\text{ cm}^{-1}$  have been found to increase, as temperature is raised, relative to intensities of other bands. Also a decrease in intensity of the band at  $933\text{ cm}^{-1}$  is observed. Since so many bands vary in intensity with temperature, a second conformer is a logical choice for the interpretation of this spectroscopic results. The less stable conformer favored at high temperatures is designated as the high energy conformer; the more stable one favored at low temperatures will be called the low energy conformer.

#### B. Intensity Considerations

The intensity of Raman bands is dependent on many factors and may be influenced by sampling methods as well as instrumental and molecular effects. New coolant and fresh sample are needed for each different temperature. Also the positions of the low



EXPLANATION OF FIGURE 17

Raman Spectrum of Liquid  $\text{CF}_2(\text{OF})_2$

Spectrometer: Spex Model 14018

Slit Setting: 150  $\mu\text{m}$

Gain: 3 K

Filter: 1.3%


Scan Speed: 1  $\text{cm}^{-1}/\text{sec}$

Chart Speed: 3  $\text{cm}/\text{min}$

Source:  $\text{Ar}^+$  Laser at 5145  $\text{\AA}$

Power: 300 mW

Temperature:  $-135^\circ\text{C}$



EXPLANATION OF FIGURE 18

Raman Spectrum of Liquid  $\text{CF}_2(\text{OF})_2$

Spectrometer: Spex Model 14018

Slit Setting: 200  $\mu\text{m}$

Gain: 3 K

Filter: 1.3%


Scan Speed: 1  $\text{cm}^{-1}/\text{sec}$

Chart Speed: 3  $\text{cm}/\text{min}$

Source:  $\text{Ar}^+$  Laser at 5145  $\text{\AA}$

Power: 300 mW

Temperature:  $-78^\circ\text{C}$



EXPLANATION OF FIGURE 19

Raman Spectrum of Liquid  $\text{CF}_2(\text{OF})_2$

Spectrometer: Spex Model 14018

Slit Setting:  $150\ \mu\text{m}$

Gain: 3 K

Filter: 1.3%


Scan Speed:  $3\ \text{cm}^{-1}/\text{sec}$

Chart Speed: 3 cm/min

Source:  $\text{Ar}^+$  Laser at  $5145\ \text{\AA}$

Power: 300 mW

Temperature:  $-40^\circ\text{C}$








Table 15. Relative Intensities<sup>a</sup> of the Bands from  
Liquid Raman Spectra  $\text{CF}_2(\text{OF})_2$

Freq. <sup>b</sup> ( $\text{cm}^{-1}$ )	Temperature ( $^{\circ}\text{C}$ )			
	-40	-78	-135	-196
108	34.8	36.8	28.0	29.5
147	4.4	6.7	4.9	6.4
243	13.2	14.1	13.0	18.0
280	16.8	16.1	12.1	14.2
304	4.0	5.0	4.9	9.2
428	6.8	8.0	6.4	12.5
434	15.2	15.4	11.3	15.3
437	37.6	36.1	32.7	41.0
524	2.8	4.0	4.6	10.9
559	6.0	5.0	3.2	4.4
602	12.4	11.4	9.0	6.8
660	42.8	44.5	41.3	49.5
692	7.6	8.0	6.6	5.8
827	4.8	4.0	2.9	2.7
837	6.4	6.7	6.4	6.8
854	33.2	33.4	32.4	32.9
860	100.0	100.0	100.0	100.0
867	64.8	50.8	35.8	20.0
874	6.8	7.7	5.8	6.4
914	17.6	17.7	15.6	12.9
933	10.4	13.0	13.3	20.0
944	34.8	30.8	22.5	16.9
956	38.8	39.1	37.3	42.7
1060	1.2	1.3	0.6	2.0
1170	3.4	3.7	3.2	3.4
1192	2.2	2.7	3.5	3.0
1207	3.2	3.0	3.8	4.1
1240	1.8	1.7	1.5	1.7

- a. The peak height of the Raman band at  $860\text{ cm}^{-1}$  was taken to be 100 for each temperature.
- b. The frequencies listed are the average values of the frequencies measured at the four temperature.



temperature cell as well as the spectrophotometer tuning may not be the same for each experiment. This means the experimental determination of absolute Raman intensities is difficult. Many corrections and calibrations are necessary before intensity values are obtained that can be used to derive molecular properties.

An approximate correction for these errors can be made by using an internal or external standard. However, with an external standard it is difficult to obtain geometrical accuracy even when using the same cell and with internal standards, the intensity of the standard may vary with environmental changes. Relative intensities can, of course, be measured (49).

We measured Raman liquid spectra at bath temperatures of  $-196$ ,  $-135$ ,  $-78$ , and  $-40^{\circ}\text{C}$  shown in Figures 16, 17, 18 and 19, respectively. Peak heights of the bands were used as a measure of the intensity of the Raman bands. Every height was normalized to a height of 100 for the most intense band which is near  $860\text{ cm}^{-1}$  at each temperature. Table 15 contains the relative intensity data measured from a 10X abscissa expansion spectrum. Thus, the relative intensities of the bands were used as a measure of the relative concentration of the indicated conformer. Figures 20 and 21 show the effect of temperature on the relative intensities in the CO and OF stretching regions.

EXPLANATION OF FIGURE 20

Effect of Temperature on the Raman Spectrum  
of Liquid  $\text{CF}_2(\text{OF})_2$  in the Range  $1000\text{-}900\text{ cm}^{-1}$

Spectrometer: Spex Model 14018

Slit Settings:  $200\text{ }\mu\text{m}$  at  $-196$  and  $-78^\circ\text{C}$

$150\text{ }\mu\text{m}$  at  $-135$  and  $-40^\circ\text{C}$

Gain: 3 K

Filter: 0.71%


Scan Speed:  $0.2\text{ cm}^{-1}/\text{sec}$

Chart Speed: 3 cm/min

Source:  $\text{Ar}^+$  Laser at  $5145\text{ }\text{\AA}$

Power: 300 mW

Temperature:  $-196$ ,  $-135$ ,  $-78$  and  $-40^\circ\text{C}$



EXPLANATION OF FIGURE 21

Effect of Temperature on the Raman Spectrum  
of Liquid  $\text{CF}_2(\text{OF})_2$  in the Range  $900\text{--}800\text{ cm}^{-1}$

Spectrometer: Spex Model 14018,

Slit Settings:  $200\text{ }\mu\text{m}$  at  $-196$  and  $-78^\circ\text{C}$

$150\text{ }\mu\text{m}$  at  $-135$  and  $-40^\circ\text{C}$

Gain: 3 K

Filter: 0.71%

Scan Speed:  $0.2\text{ cm}^{-1}/\text{sec}$

Chart Speed: 3 cm/min

Source:  $\text{Ar}^+$  Laser at  $5145\text{ }\text{\AA}$

Power: 300 mW

Temperature:  $-196$ ,  $-135$ ,  $178$  and  $-40^\circ\text{C}$




Table 16. Summary of Tentative Assignment for a 2 Conformer Mixture

Low energy conformer			High energy conformer		
Raman liq. <sup>a</sup>	Raman gas <sup>b</sup>	IR gas	Raman liq. <sup>a</sup>	Raman gas <sup>b</sup>	IR gas
	1287.5	1275		1287.5	1275
1240	1246.1	1247	1240	1246.1	1247
1207	1215.6	1207	1207	1215.6	1207
956	956.9	955	944	943.8	942
933	931.0	933Q	914	915.4	916Q
			874		
860	854.9	857	867	863.1	866
854		~850			
837		836.5	827		831
660	657.7	658.5Q	692	687.3	692Q
602 or No	601.9 or No	598 or No	602	601.9	603
524	525?	522.5Q	559	557.3	555Q
437	No?	No, ~440?	434	432.5	No
428 or No	No?	426	428	No?	426
304	No?	300Q or No	291	No?	300Q
243	241.1	~240	280	278.9	~280
147			147		
108			108		

- a. The liquid Raman frequencies listed are the average values of the frequencies measured at the four different temperatures.
- b. These frequencies are from reference (19) where the data are more complete than ours. Our gas phase Raman spectra in Figure 15 differ in frequency for the most prominent bands in the  $800\text{--}1000\text{ cm}^{-1}$  region where we find 958, 945, 867 and  $860\text{ cm}^{-1}$  rather than the values 956.9, 943.8, 863.1 and  $854.9\text{ cm}^{-1}$  from reference (19) listed above.

From Figures 20 and 21 it can be seen that the relative intensities of the bands at 956, 933, 860 and 854  $\text{cm}^{-1}$  decrease when the temperature increases, these bands are classified as belonging to the low energy conformer. The reverse is true for the bands at 944, 914 and 867  $\text{cm}^{-1}$ , and they are classified as belonging to the high energy conformer.

A summary of tentative assignments for a 2 conformer or rotational isomer mixture made by taking the available spectroscopic data for comparison is listed in Table 16.

#### C. Energy Difference between Rotational Isomers

We have concluded that temperature effects in the liquid state could be explained by the presence of the two rotational isomers: high energy conformer or isomer and low energy conformer or isomer.

As seen in Figure 20, relative intensities of the Raman bands at 944 and 956  $\text{cm}^{-1}$  vary with temperature, indicating definitely that they belong to different rotational isomers. Since the Raman bands at 944 and 956  $\text{cm}^{-1}$  are now assigned to the two different conformers, we can determine the energy difference between the isomers from the temperature dependence of the Raman intensity ratios. It was shown that there seems to be no significant difference in the results obtained whether peak height intensities or band area intensities are used for energy difference calculations (50).

Although the Raman bands at 944 and 956  $\text{cm}^{-1}$  partially overlap each other, the peak heights of these two bands were used as a measure of the concentration of the indicated isomer based on the above approximation.

The variations are given as follows:

$$-196^{\circ}\text{C} = 77^{\circ}\text{K} \quad -135^{\circ}\text{C} = 138^{\circ}\text{K} \quad -78^{\circ}\text{C} = 195^{\circ}\text{K}$$

$$\frac{I_{956}}{I_{944}} = \quad 2.53 \quad 1.66 \quad 1.27$$

The energy difference  $\Delta E = E(\text{high energy}) - E(\text{low energy})$  was evaluated by the equation (51):

$$\left( \frac{I_{956}}{I_{944}} \right)_{T_1} \bigg/ \left( \frac{I_{956}}{I_{944}} \right)_{T_2} = \exp \left[ -\frac{\Delta E}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right) \right]$$

The values calculated for  $T_1 = 77^{\circ}\text{K}$  and  $T_2 = 138^{\circ}\text{K}$ ,  $T_1 = 138^{\circ}\text{K}$  and  $T_2 = 195^{\circ}\text{K}$ , and  $T_1 = 77^{\circ}\text{K}$  and  $T_2 = 195^{\circ}\text{K}$  were 146, 251, and 174 cal/mole, respectively. Since the data at  $138^{\circ}\text{K}$  were recorded with a different slit setting and since  $77^{\circ}\text{K}$  and  $195^{\circ}\text{K}$  are the most widely separated temperatures below the boiling point of  $\text{CF}_2(\text{OF})_2$  ( $-64^{\circ}\text{C} = 209^{\circ}\text{K}$ ),  $\Delta E = 174$  cal/mole is probably the most reliable value. The data from  $-40^{\circ}\text{C}$  was not used for fear of possible vaporization during the recording of the spectrum. An analogous calculation using the 867 and 860  $\text{cm}^{-1}$  bands gave 202, 326, and 235 cal/mole, but these values should be less reliable due to more serious overlap between the 867 and 860  $\text{cm}^{-1}$  bands than between the



944 and 956  $\text{cm}^{-1}$  bands (see Figures 20 and 21). Since a lot of approximations are involved in the calculation, it can only be concluded that energy difference between the two isomers must be quite small and probably is of the order of 100-300 cal/mole.

#### D. Suggestions for Future Work

The experimental results presented here demonstrate that the liquid Raman data can be interpreted in terms of an equilibrium mixture of two rotational isomers over the liquid range of  $\text{CF}_2(\text{OF})_2$  (b. p.  $-64^\circ\text{C}$ ) down to near  $-196^\circ\text{C}$ . Other measurements that might be informative include the following: infrared and Raman spectra of the pure solid and the solid isolated in an argon matrix, very low temperature F-19 NMR spectra of the liquid, Raman spectra of the liquid with accurate depolarization ratio measurements, and microwave spectra of the gas phase.

If a crystalline solid phase will form below  $-196^\circ\text{C}$  or if a solid may be isolated in an argon matrix, there may be so little of the high energy isomer present that it will make a negligible contribution to the IR and Raman spectra. Then the fundamentals due to the low energy isomer might be conclusively identified.

If the rate of rotation from one isomer to the other can be slowed sufficiently the F-19 NMR signals will be split into signals for each isomer and the chemical shifts and coupling

constants of each isomer might be obtained. These data might assist in establishing the structure of each isomer.

If one could correctly identify all the depolarized bands in the Raman spectrum of the liquid, then the data in this work would indicate how many depolarized bands each isomer has. Such information might allow one or both isomers to be assigned to the proper point group.

Recently, microwave spectroscopy has been used to identify rotational isomers (52). The gas phase Raman spectrum of  $\text{CF}_2(\text{OF})_2$  near room temperature in Figure 15 suggests that the two isomers are present in about equal concentrations since the gas phase analogues of the 867 and 860  $\text{cm}^{-1}$  bands in the liquid have about the same intensity. Consequently, it might be possible to assign lines to both isomers and obtain structural data for both isomers.

## LITERATURE CITED

1. I. W. Levin and R. A. R. Pearce, in Vibrational Spectra and Structure, Vol. 4, (J. R. Durig, ed.), Elsevier Scientific, Amsterdam, 1975, p. 101.
2. C. J. Hoffman, Chem. Rev., 64, 91 (1964).
3. C. J. Schack and W. Maya, J. Am. Chem. Soc., 91, 2902 (1969).
4. P. M. Wilt and E. A. Jones, J. Inorg. Nucl. Chem., 30, 2933 (1968).
5. R. R. Smardzewski and W. B. Fox, J. Fluoro. Chem., 6, 417 (1975).
6. P. Buckley and J. P. Weber, Can. J. Chem., 52, 942 (1974).
7. C. J. Marsden, D. D. DesMarteau, and L. S. Bartell, Inorg. Chem., 16, 2359 (1977).
8. Ajit S. Manocha, M. S. Thesis, Kansas State University, Manhattan, Kansas (1978).
9. E. B. Wilson, Jr., J. C. Decius, and P. C. Cross, Molecular Vibrations, McGraw-Hill, New York, 1955.
10. J. H. Schachtschneider, "Vibrational Analysis of Polyatomic Molecules. V and VI", Tech. Rept. Nos. 231-64 and 57-65, respectively, Shell Development Co., Houston, Texas.
11. K. B. Kellog and G. H. Cady, J. Am. Chem. Soc., 70, 3986 (1948).
12. R. T. Lagemann, E. A. Jones, and P. J. H. Woltz, J. Chem. Phys., 20, 1768 (1952).
13. F. P. Diodati and L. S. Bartell, J. Mol. Structure, 8, 395 (1971).
14. D. D. DesMarteau, Kansas State University, private communication.
15. D. E. Gould, L. R. Anderson, D. E. Young, and W. B. Fox, Chem. Commun., 1564 (1968).

16. M. Lustig, A. R. Pitochelli, and J. K. Ruff, J. Am. Chem. Soc., 89, 2841 (1967).
17. I. U. P. A. C., Tables of Wavenumbers for the Calibration of Infrared Spectrometers, Butterworth, Washington, D. C. 1961.
18. C. W. Brown, A. G. Hopkins, and F. P. Daly, Appl. Spectry., 28, 194 (1974).
19. R. R. Smardzewski and W. B. Fox, U. S. Naval Research Laboratory, Washington, D. C. 20375, private communication to Dr. D. D. DesMarteau.
20. J. C. Decius, J. Chem. Phys., 17, 1315 (1949).
21. John R. Ferraro and Joseph S. Ziomek, Introductory Group Theory, 2nd Ed. Plenum Press, New York, 1975.
22. J. H. Schachtschneider and R. G. Snyder, Spectrochim. Acta, 19, 117 (1963).
23. D. E. Mann, T. Shimanouchi, J. H. Meal, and L. Fano, J. Chem. Phys., 27, 43 (1957).
24. W. T. King, I. M. Mills, and B. L. Crawford, Jr., J. Chem. Phys., 27, 455 (1957).
25. J. Overend and J. R. Scherer, J. Chem. Phys., 32, 1289 (1960).
26. T. Shimanouchi and I. Suzuki, J. Chem. Phys., 42, 296 (1965).
27. L. Pierce, N. Di Cianni, and R. Jackson, J. Chem. Phys., 38, 730 (1963).
28. Y. Morino and S. Saito, J. Mol. Spectry., 19, 435 (1966).
29. P. N. Noble and G. C. Pimentel, Spectrochim. Acta, 24A, 797 (1968).
30. J. F. Ogilvie, Canad. J. Spectry., 19, 171 (1974).
31. M. M. Rochkind and G. C. Pimentel, J. Chem. Phys., 42, 1361 (1965).
32. B. Beagley, A. H. Clark, and D. W. J. Cruickshank, Chem. Comm., 458 (1966).

33. G. E. Herberich, R. H. Jackson, and D. J. Millen, J. Chem. Soc. (A), 336 (1966).
34. K. O. Christe, C. J. Schack, and E. C. Curtis, Inorg. Chem., 10, 1589 (1971).
35. J. D. Witt and R. M. Hammaker, J. Chem. Phys., 58, 303 (1973).
36. R. N. Dixon, J. Chem. Phys., 31, 258 (1959).
37. E. L. Saier, L. R. Cousins, and M. Basila, J. Phys. Chem., 66, 232 (1962).
38. W. G. Fateley, I. Matsubara, and R. E. Witkowski, Spectrochim. Acta, 20, 1461 (1964).
39. T. Shimanouchi, Tables of Molecular Vibrational Frequencies, Consolidated Volume, NSRDS-NBS-39 (National Bureau of Standards, Washington, D. C., 1972), p. 22.
40. G. M. Begun, and W. H. Fletcher and D. F. Smith, J. Chem. Phys., 42, 2236 (1965).
41. H. Selig, H. H. Claassen, and J. H. Holloway, J. Chem. Phys., 52, 3517 (1970).
42. W. G. Fateley and F. A. Miller, Spectrochim. Acta, 17, 857 (1961).
43. W. G. Fateley and F. A. Miller, Spectrochim. Acta, 18, 977 (1962).
44. W. G. Fateley, F. A. Miller, and R. E. Witkowski, Technical Report AFML-TR-66-408, Air Force Materials Laboratory, Wright-Patterson Air Force Base, Ohio 45433.
45. P. G. Thompson, "Oxygen Fluorides and Hypofluorites", The Fluorine Symposium of the Inorganic Division of the American Chemical Society, Ann Arbor, Mich., June 27, 1966.
46. P. Thompson, J. Am. Chem. Soc., 89, 1811 (1967).
47. F. A. Hohorst and J. M. Shreeve, J. Am. Chem. Soc., 89, 1809 (1967).
48. R. W. Mitchell and J. A. Merrit, J. Mol. Spectry., 24, 128 (1967).

49. P. J. D. Park, R. A. Pethrick and B. H. Thomas, in Internal Rotation in Molecules, (W. J. Orville Thomas ed.), John Wiley & Sons, London, 1974, p.83.
50. H. J. Bernstein, J. Chem. Phys., 17, 258 (1949).
51. A. Langeseth and H. J. Bernstein, J. Chem. Phys., 8, 410 (1940).
52. N. S. True and R. K. Bohn, J. Phys. Chem., 81, 1667, 1671 (1977).
53. A. Müller and B. Krebs, J. Mol. Spectroscopy, 24, 180 (1967); B. Krebs, A. Müller, and A. Fadini, J. Mol. Spectroscopy, 24, 198 (1967); J. L. Duncan and I. M. Mills, Spectrochim Acta, 20, 1089 (1964).
54. A. Ruoff, H. Burger, and S. Biedermann, Spectrochim. Acta, 27A, 1359 (1971); R. W. Kirk and P. M. Wilt, J. Mole. Spectroscopy, 58, 102 (1975).

## ACKNOWLEDGMENT

A considerable portion of one's destiny is molded by those people with whom he is associated. While one must accept the ultimate responsibility for the realization of his objectives, it is obvious that they can not be achieved without the assistance and encouragement of others.

In appreciation of this, the author would like to extend his sincere gratitude to Professor R. M. Hammaker for his invaluable contributions to this work. Also for his patience, valuable discussion and all his help during the course of this study, the author is deeply appreciative.

The author is deeply indebted to Professors W. G. Fateley and D. D. DesMarteau for their incomparable guidance and assistance over the year when Dr. Hammaker was abroad. Also the help from the Flourine Chemistry Laboratory of Kansas State University deserves the author's special thanks.

Last but not least, the author wishes to convey special praise to his wife, Fen. Without her love and understanding, none of this work would have been accomplished.

## VITA

The author was born on February 21, 1950, in Yun-Lin, Taiwan, Republic of China. He obtained his B.S. degree in Chemistry from National Taiwan University in June, 1972. He began graduate study at Kansas State University in August, 1974. He is a member of Phi Lambda Upsilon.



STUDIES IN VIBRATIONAL SPECTROSCOPY

- I. NORMAL COORDINATE ANALYSIS OF  $\text{CF}_3\text{OF}$  AND  $\text{CF}_3\text{OCl}$
- II. RAMAN SPECTRA OF  $\text{CF}_2(\text{OF})_2$  AS A FUNCTION OF TEMPERATURE

by

JENG-CHUNG KUO

B.S., National Taiwan University, 1972

---

AN ABSTRACT OF A MASTER'S THESIS

Submitted in partial fulfillment of the  
requirements for the degree

MASTER OF SCIENCE

Department of Chemistry

KANSAS STATE UNIVERSITY  
Manhattan, Kansas

1978

## I. NORMAL COORDINATE ANALYSIS OF $\text{CF}_3\text{OF}$ AND $\text{CF}_3\text{OCl}$

Raman and infrared data previously obtained in this laboratory are used to perform a normal coordinate analysis on  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$ . All infrared and Raman bands below  $1400\text{ cm}^{-1}$  have been assigned under  $\text{C}_s$  symmetry. Both the fundamental and the first overtone of the  $\text{CF}_3$  torsion in both  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  have been observed. The two  $\text{CF}_3$  rocking modes of both  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$  appear to be well separated in frequency. The barriers to internal rotation of the  $\text{CF}_3$  group are 4.4 and 4.2 kcal/mole in  $\text{CF}_3\text{OF}$  and  $\text{CF}_3\text{OCl}$ , respectively. An unexpected band near  $548\text{ cm}^{-1}$  in the Raman spectrum of liquid  $\text{CF}_3\text{OCl}$  near  $77^\circ\text{K}$  is tentatively assigned to the stretching of the chlorine to chlorine bond of molecular chlorine in a complex with  $\text{CF}_3\text{OCl}$ .

## II. RAMAN SPECTRA OF $\text{CF}_2(\text{OF})_2$ AS A FUNCTION OF TEMPERATURE

Raman and infrared data have been obtained for  $\text{CF}_2(\text{OF})_2$ . The large number of bands in the Raman spectrum of the liquid have been interpreted in terms of a mixture of two rotational isomers or conformers. Raman spectra of liquid  $\text{CF}_2(\text{OF})_2$  at  $-196$ ,  $-135$ ,  $-78$ , and  $-40^\circ\text{C}$  have been used to assign the observed Raman and infrared bands to a high energy conformer and low energy conformer. Using peak heights of strong Raman bands between  $800$  and  $1000\text{ cm}^{-1}$ , the energy separation between the two conformers is estimated to be in the range  $100$  to  $300\text{ cal/mole}$ .